

(FAPP) Infinity Does Macroscopic Irreversibility From Microscopic Reversibility

Karl Svozil^{1,*}

¹*Institute for Theoretical Physics, TU Wien, Wiedner Hauptstraße 8-10/136, 1040 Vienna, Austria*

(Dated: April 29, 2025)

Infinity is central to deriving macroscopic irreversibility from reversible microscopic laws across mathematics, theoretical computer science and physics. In analysis, infinite processes—such as Dedekind cuts and Cauchy sequences—construct real numbers as equivalence classes of rational approximations, bridging discrete rationals to the continuous real line. In quantum mechanics, infinite tensor products model nested measurements, where sectorization partitions the Hilbert space into equivalence classes, reconciling unitary evolution with wavefunction collapse. In statistical mechanics, macrostates emerge as equivalence classes of microstates sharing identical macroscopic properties, providing the statistical basis for thermodynamic irreversibility despite reversible dynamics. Equivalence relations formalize For-All-Practical-Purposes (FAPP) indistinguishability, reflecting operational limits on precision and observation. Together, these examples reveal a unified framework where infinity and equivalence underpin emergent macroscopic behavior from microscopic reversibility.

Keywords: infinity, FAPP, Specker sequence, Chaitin's Omega, halting probability

I. FROM RATIONALS TO REALS: THE ROLE OF INFINITY

The construction of the real numbers from the rational numbers is a fundamental topic in mathematical analysis, highlighting the necessity of infinite processes. The rational numbers, denoted by \mathbb{Q} , are countable and dense in the real numbers, but they are incomplete. This incompleteness arises because there exist ‘gaps’ in \mathbb{Q} that correspond to irrational numbers. To fill these gaps and construct the real numbers, \mathbb{R} , mathematicians employ infinite methods, such as continued fractions. Two prominent approaches are *Dedekind cuts* and *Cauchy sequences*, both of which rely on the concept of infinity. The discussion will explore methods employing infinite means that transcend from rational to irrational numbers, then progress through Specker sequences to uncomputable numbers, and finally examine Omega sequences leading to algorithmically incompressible random reals. At this point, concerns about the physical operationality of these infinite means will be set aside, with the issue revisited later in the discussion.

A. Dedekind Cuts

A Dedekind cut partitions the rational numbers into two non-empty sets A and B such that every element of A is less than every element of B , and A contains no greatest element. The cut represents a real number, which may be rational or irrational. For example, the cut corresponding to $\sqrt{2}$ is defined by:

$$A = \{x \in \mathbb{Q} \mid x^2 < 2\}, \quad B = \{x \in \mathbb{Q} \mid x^2 > 2\}.$$

This construction inherently involves an infinite sets A and B . The completeness of the real continuum is embodied by the property that every such cut corresponds to a unique real

number, effectively filling the ‘irrational gaps between’ rationals. This construction vividly illustrates that the limit of a sequence—often an irrational number—can be captured only through an infinite process corresponding to the infinite sets A and B .

Similarly, surreal numbers, introduced by Conway [1] and explored in a mathematical dialogue by Knuth [2], are constructed recursively as equivalence classes of pairs of sets of surreal numbers, subject to the condition that every element of the first set is less than every element of the second set. The construction begins with the empty set. At each stage, new numbers are defined as $\{L \mid R\}$, where L and R are sets of previously constructed numbers, provided that every member of L is less than every member of R . This Dedekind cut-like procedure, iterated transfininitely and allowing L and R to be infinite, produces not only all standard real numbers but also a vast continuum of infinite and infinitesimal numbers. Thus, from the initial void—the empty set $\{\mid\}$ identified with the number 0—this infinite process generates a comprehensive universe of numbers, truly *ex nihilo omnia* (everything out of nothing).

B. Cauchy Sequences

Another method to construct the real numbers is through Cauchy sequences of rational numbers. A Cauchy sequence $(x_n)_{n=1}^{\infty}$ is a sequence whose elements become arbitrarily close to each other as the sequence progresses. Formally, for every $\varepsilon > 0$, there exists an integer N such that for all $m, n \geq N$, $|x_m - x_n| < \varepsilon$. The real numbers are then defined as equivalence classes of Cauchy sequences, where two sequences are equivalent if their difference converges to zero. This process also relies on infinity, as the convergence of the sequence is an infinite phenomenon.

* karl.svozil@tuwien.ac.at; <http://tph.tuwien.ac.at/~svozil>

C. Infinite Decimal Expansions

A more familiar representation is that of *infinite decimal expansions*. Any real number can be expressed as an infinite sequence of digits, $x_0.x_1x_2x_3\dots$, which in turn can be viewed as an infinite sum. This representation not only emphasizes the necessity of an infinite process but also shows how numbers that cannot be finitely represented (such as irrational numbers) naturally arise from the completion of an endless procedure.

D. Cantor's Diagonalization and Irrational Numbers

Cantor's diagonalization argument is a powerful tool that demonstrates the uncountability of the real numbers and provides a method to construct irrational numbers from rationals through an infinite process. Consider an enumeration of all rational numbers in the interval $[0, 1]$, say r_1, r_2, r_3, \dots . Each rational number r_i can be expressed as an infinite decimal expansion. By constructing a new number x whose n -th decimal digit differs from the n -th decimal digit of r_n , we ensure that x is distinct from every rational number in the list. For instance, if the n -th digit of r_n is d_n , define the n -th digit of x as $d_n + 1 \bmod 10$. The resulting number x is irrational, as it cannot correspond to any rational number in the enumeration. This construction explicitly relies on an infinite process that 'constructs' an irrational number [3, 4].

E. No Continua Without Infinite Means

The transition from the rational numbers to the real numbers—whether through Dedekind cuts, Cauchy sequences, or infinite decimal expansions—as well as Cantor's diagonalization argument necessitates the use of infinite means. These Zeno-type constructions underscore the indispensable role of infinity in bridging the gap between the countable realm of \mathbb{Q} and the uncountable continuum of \mathbb{R} : No finite procedure that starts with a finite set of rational numbers and uses only a finite number of operations can produce an irrational number.

Whether the infinities inherently present in (classical) continua can be put to any operational physical use remains an open question [5]. Suffice it to say that the assumption of continua, as well as the selection of one of their elements via the axiom of choice, is a key ingredient in the apparent oxymoron that is the widely used term deterministic chaos.

Noson Yanofsky has noted that the procedural approach used here to generate the continuum and other mathematical entities, such as irrational or uncomputable numbers, including through methods like diagonalization, could be criticized. The criticism stems from the view that tools like Dedekind cuts and Cauchy sequences describe or represent numbers rather than actually constructing them [6]. However, while this raises a relevant metamathematical concern, it ultimately hinges on a matter of philosophical perspective.

F. Specker Sequences and the Role of Infinity

Just as infinity plays an indispensable role in the transition from rational to irrational numbers and in the conceptualization of mathematical continua, *Specker sequences* provide a profound illustration of how infinity can lead us from the computable to the uncomputable, thereby selecting a subset of irrationals by tightening criteria. Almost all reals are of this type.

A Specker sequence is a computable, monotonically increasing, bounded sequence of rational numbers whose limit is an uncomputable real number [7, 8]. Formally, a sequence $(a_n)_{n=1}^{\infty}$ is a Specker sequence if:

1. Each a_n is a computable rational number
2. The sequence is strictly increasing: $a_n < a_{n+1}$ for all n
3. The sequence is bounded above: there exists $L \in \mathbb{Q}$ such that $a_n < L$ for all n
4. The limit $\lim_{n \rightarrow \infty} a_n$ is not a computable number

The existence of such sequences demonstrates that the infinite completion of even well-behaved, computable objects can yield entities beyond algorithmic reach. The essence of Specker's construction is to encode an undecidable property into the convergence behavior of the sequence. Although each term a_n is produced by a finite, effective algorithm, the process of converging to L is intrinsically infinite—any attempt to specify a convergence criterion would require solving a problem that is uncomputable—indeed, to quote an early, informal intuition by Paul Ehrenfest, such a convergence criterion “grows beyond any specifiable size” [9]. In this way, the limit L becomes an uncomputable real number even though it is the limit of a computable (recursive) sequence.

G. Chaitin's Omega as the Ultimate Specker Sequence

Perhaps the most profound example of a limit of a Specker sequence is Chaitin's Omega (Ω), often called the ‘halting probability’ [10]. This number represents the probability that a randomly constructed self-delimiting program will halt when run on a universal Turing machine.

Chaitin's Ω can be expressed as:

$$\Omega = \sum_{p \text{ halts}} 2^{-|p|}$$

where the sum is taken over all self-delimiting programs p that halt, and $|p|$ denotes the length of program p in bits.

Ω can be approximated by rational numbers:

$$\Omega_n = \sum_{\substack{p \text{ halts within} \\ n \text{ steps}}} 2^{-|p|}$$

The sequence $(\Omega_n)_{n=1}^{\infty}$ is a Specker sequence—each Ω_n is computable for ‘small’ n [11], the sequence is monotonically

increasing, bounded above by 1, yet its limit Ω is uncomputable. The uncomputable nature of Ω stems from the fact that knowledge of its binary expansion would allow us to solve the Halting Problem, which is provable impossible. Each additional bit of precision in Ω encodes the solution to increasingly complex instances of the Halting Problem. There does not exist any computable convergence criterion: just as for computing the n th bit of a Busy Beaver function, the time to compute those instances Ω_n outgrows any computable function of n [12].

All of the above reveals a fundamental qualitative shift at infinity—one that goes beyond mere quantitative change: Infinity generates fundamentally new mathematical objects. The rational numbers, all of which are computable, give rise through infinite processes to real numbers that no algorithm can fully capture.

The transition from the finite to the infinite marks a profound divide between what is algorithmically accessible and what remains beyond reach. While we can approximate Ω arbitrarily closely using computable methods, we can never compute it exactly. Specker sequences thus demonstrate that infinity is not merely a convenient mathematical abstraction but a necessary concept that marks the boundary between the computable and the uncomputable, between what can be algorithmically constructed and what can only be defined through infinite convergence. Indeed, despite random reals [13] constituting almost all irrational numbers, locating specific instances through computational, finite, or physically operational means remains provably impossible [14].

Specker sequences and Chaitin's Omega are not defined using equivalence classes in their original formulations. But they are related to equivalence classes in how they can be introduced: They are indirectly tied through the Cauchy sequence construction of real numbers, where their limits are equivalence classes.

II. INFINITE TENSOR PRODUCTS AND THE QUANTUM MEASUREMENT PROBLEM

Infinite tensor products, when interpreted as infinite chains of nested measurements, provide a compelling framework for addressing the quantum measurement problem. By introducing disruptions to unitary equivalence through sectorization and factorization, this approach offers a potential reconciliation between the unitary evolution of quantum systems and the apparent collapse of the wavefunction during measurement.

The quantum measurement problem remains one of the most profound challenges in quantum mechanics, arising from the apparent inconsistency between two fundamental processes identified by von Neumann in 1932 [15–17]. These processes are:

Process 1: The discontinuous, probabilistic change in a quantum state upon measurement. For a system in a superposition $\psi = \sum_i c_i \phi_i$, observing a quantity with eigenstates ϕ_1, ϕ_2, \dots collapses the state to ϕ_j with probability $|c_j|^2$.

Process 2: The continuous, deterministic evolution of an isolated system's state according to the Schrödinger equation, $\partial\psi/\partial t = U\psi$, where U is a unitary operator.

The crux of the measurement problem is whether the unitary evolution (Process 2) can fully account for the collapse observed in measurements (Process 1), or if an additional mechanism is required. This section explores the use of infinite tensor products, interpreted as infinite nestings of Wigner's friend scenarios, as a potential resolution to this problem.

A. Infinite Tensor Products in Nested Measurement Scenarios

A promising approach to addressing the measurement problem involves infinite tensor products, which model an infinite sequence of observers, each measuring the system observed by the previous observer. This setup is reminiscent of Wigner's friend thought experiments, where the act of measurement is recursively applied. Unlike finite tensor products, infinite tensor products can disrupt unitary equivalence through mechanisms such as sectorization and factorization, potentially providing a bridge between unitary evolution and the apparent collapse of the wavefunction.

B. The Von Neumann-Landau Measurement Scheme

In the von Neumann-Landau framework, the measurement process is modeled by the interaction between an object and a measurement apparatus. The object is prepared in a state $|\Psi\rangle = \sum_{i=1}^n a_i |\Psi_i\rangle$, which is a superposition relative to the measurement basis. The measurement apparatus is represented by another state $|\phi\rangle = \sum_{j=1}^m b_j |\phi_j\rangle$. Upon interaction, the combined state of the object and apparatus becomes:

$$|\Psi\rangle = \sum_{i,j=1}^n c_{ij} |\Psi_i\rangle \otimes |\phi_j\rangle,$$

where the coefficients c_{ij} cannot be factorized, indicating entanglement between the object and the apparatus.

While this scheme is straightforward for finite systems, extending it to an infinite chain of measurements—where each measurement is itself measured by another observer, ad infinitum—requires the use of infinite tensor products. This extension is mathematically non-trivial and was first rigorously studied by von Neumann in 1939 [18].

The construction of the infinite tensor product space proceeds as follows:

1. Begin with elementary tensors of the form $\otimes_{n=1}^{\infty} |k_n\rangle$.
2. Define the inner product between two elementary tensors as:

$$\left\langle \left(\bigotimes_{n=1}^{\infty} |k_n\rangle \right) \left| \bigotimes_{n=1}^{\infty} |l_n\rangle \right. \right\rangle = \prod_{n=1}^{\infty} \langle k_n | l_n \rangle,$$

provided the product converges; otherwise, it is zero.

3. Consider finite linear combinations of these elementary tensors:

$$\sum_i c_i \bigotimes_{n=1}^{\infty} |k_n^{(i)}\rangle,$$

where c_i are complex coefficients and $|k_n^{(i)}\rangle$ are basis vectors.

4. Obtain the complete Hilbert space $\bigotimes_{n=1}^{\infty} \mathcal{H}_n$ by taking the closure of the space of finite linear combinations.

This construction introduces several challenges that must be addressed to fully understand its implications for the measurement problem.

C. Challenges with Infinite Tensor Products

1. Cardinality

As pointed out by von Neumann in 1939 [18], a fundamental issue with infinite tensor products is the uncountable cardinality of the resulting space. Just as the real numbers cannot be enumerated by a countable set, the infinite tensor product space cannot be spanned by a countable basis. Such generalizations involve nonseparable Hilbert spaces and higher set-theoretical powers of their orthonormal bases, thereby spoiling unitary equivalence with (in)finite-dimensional separable Hilbert spaces.

The interval $[0, 1)$ can be represented in binary form as

$$\{0.x_1x_2x_3\dots \mid x_i \in \{0, 1\} \text{ for all } i \in \mathbb{N}\}.$$

Here, each x_i is a binary digit (0 or 1), and the sequence extends indefinitely. This set of all infinite binary sequences is uncountable, with cardinality

$$\#\{0, 1\}^{\mathbb{N}} = 2^{\aleph_0},$$

which, by Cantor's diagonal argument mentioned earlier, is strictly larger than the cardinality \aleph_0 of the natural numbers.

In close analogy, consider an infinite sequence of qubits, where each qubit is a two-state quantum system with basis states $|0\rangle$ and $|1\rangle$. A product state in the infinite tensor product is written as

$$|x_1x_2x_3\dots\rangle = |x_1\rangle \otimes |x_2\rangle \otimes |x_3\rangle \otimes \dots,$$

with $x_i \in \{0, 1\}$ for all $i \in \mathbb{N}$. The collection of all such product states corresponds exactly to the set of infinite binary sequences, hence its cardinality is also 2^{\aleph_0} .

A denumerable set of product states is any countable subset of the infinite tensor product states. For instance, the set

$$\{|000\dots\rangle, |100\dots\rangle, |010\dots\rangle, \dots\}$$

can be put into a one-to-one correspondence with the natural numbers \mathbb{N} and thus has cardinality \aleph_0 . Clearly, $\aleph_0 < 2^{\aleph_0}$.

A Hilbert space is *separable* if it has a countable (finite or infinite) orthonormal basis; otherwise it is called nonseparable. Two Hilbert spaces \mathcal{H}_1 and \mathcal{H}_2 are unitarily equivalent if and only if they have the same *dimension* (that is, their orthonormal bases share the same cardinality). More explicitly, if $|e_i\rangle$ and $|f_i\rangle$ are denumerable orthonormal bases for separable \mathcal{H}_1 and \mathcal{H}_2 , respectively, then the unitary operator is given by

$$U = \sum_i |f_i\rangle\langle e_i|.$$

In our scenario, the full Hilbert space of the infinite tensor product space is nonseparable, as it has an orthonormal basis consisting of product states with cardinality 2^{\aleph_0} . Any candidate unitary operator mapping a countable (denumerable) subset of product states (with cardinality \aleph_0) to the full set must preserve the inner-product structure and be surjective onto the basis. However, since a unitary map must preserve the cardinality of an orthonormal basis, and we have $\aleph_0 < 2^{\aleph_0}$, no such unitary operator can exist that maps a countable subset onto the full uncountable basis.

2. Inner Product and Orthogonality

Another significant challenge arises in defining the inner product for infinite tensor products. For two states $|\Psi\rangle = \bigotimes_{i=1}^{\infty} |x_i\rangle$ and $|\Phi\rangle = \bigotimes_{i=1}^{\infty} |y_i\rangle$, the inner product is given by:

$$\langle \Psi | \Phi \rangle = \prod_{i=1}^{\infty} \langle x_i | y_i \rangle.$$

If each $\langle x_i | y_i \rangle = 1 - \varepsilon_i$ with $0 < \varepsilon_i \ll 1$, and if the series $\sum_{i=1}^{\infty} \varepsilon_i$ diverges, then:

$$\prod_{i=1}^{\infty} (1 - \varepsilon_i) \approx \exp \left(- \sum_{i=1}^{\infty} \varepsilon_i \right) \rightarrow 0.$$

This implies that states which are only slightly different across infinitely many components can have an inner product that approaches zero, making them effectively orthogonal. Moreover, if the states differ in even a single component such that $\langle x_k | y_k \rangle = 0$ for some k , the entire inner product becomes zero, regardless of the similarity in other components. This behavior disrupts traditional notions of orthogonality and complicates the interpretation of measurement outcomes.

D. Sectorization as a Solution

To address these issues, von Neumann proposed partitioning the infinite tensor product space into disjoint 'regions' or *sectors*—equivalence classes of states that are 'close' to each other in a specific sense [18]. Two states $|\Psi\rangle$ and $|\Phi\rangle$ are considered to be in the same sector if:

$$\sum_{i=1}^{\infty} (1 - |\langle x_i | y_i \rangle|) < \infty.$$

This condition ensures that the states differ significantly in only finitely many components. These sectors can be thought of as corresponding to distinct macroscopic or classical outcomes, potentially offering a way to interpret measurement results within the framework of unitary evolution [19–22].

E. Factorization and Unitary Equivalence

A further opportunity arises from the entanglement of infinite components, which can lead to different types of factors (e.g., type I, II, or III in von Neumann algebra classification) that are not unitarily equivalent. This lack of unitary equivalence suggests a mechanism by which the infinite tensor product space can accommodate irreversible processes, such as those observed in quantum measurements.

F. Role of equivalence classes

In sectorization, equivalence classes are employed to partition the infinite tensor product space into distinct sectors, each comprising states that are equivalent modulo differences in only finitely many components. This classification is pivotal for associating each sector with a specific, classical measurement outcome, thereby offering a framework to reconcile the continuous, unitary evolution of quantum systems with the discrete nature of observed measurement results. Furthermore, the lack of unitary equivalence between different sectors—stemming from the factorization of the space—underscores the critical role of equivalence classes in establishing the irreversibility characteristic of the quantum measurement process. By defining these equivalence classes, sectorization simplifies the handling of complex quantum systems and provides insight into the transition from quantum superpositions to definite classical states.

III. INFINITE PRECISION MICROSTATES AND THE EMERGENCE OF MACROSCOPIC IRREVERSIBILITY

A common starting point in statistical physics is to describe an isolated many-particle system by specifying its *microstate* with infinite precision. In principle, if every particle's position and momentum were known exactly, the time-reversible microscopic laws (that is, Newtonian or unitary quantum dynamics) imply that every evolution has a time-reversed twin. This observation is at the heart of Loschmidt's *Umkehrereinwand* (reversal objection) [23]: If one were able to precisely reverse the velocities of all particles, then every macroscopic process (such as the free expansion of a gas) would be exactly reversible. In other words, entropy would remain constant with infinite precision.

In an extreme scenario where a hidden entity, such as Maxwell's demon, manipulates a system at the microphysical level with infinite precision—unbeknownst to observers who are limited to finite precision macroscopic measurements—the demon could orchestrate processes like the spontaneous

unmixing of two previously mixed gases. This would make entropy appear to decrease from the observers' macroscopic perspective, creating the illusion of a contradiction with the second law of thermodynamics, which states that entropy in an isolated system cannot decrease over time.

However, concepts such as physical means and demons manipulating microstates with infinite precision is an idealized notion. In reality, our ability to specify, measure, or manipulate microscopic degrees of freedom in any physical system is operationally limited. These practical constraints mean that microstates can only be defined with *finite* precision. Consequently, when attempting to reverse a system's evolution, the unavoidable small uncertainties are amplified through the system's complex (often chaotic) dynamics.

Moreover, the concept of *means-relative reversibility* emphasizes that while the microscopic laws are symmetric, the notion of a reversible process depends on the precision and the scale at which the state is defined. Maxwell's pragmatic approach—“avoiding all personal inquiries [[about individual molecules]] which would only get me into trouble” [24, 25]—illustrates that the coarse-grained description relevant for thermodynamics deliberately sidesteps the need for infinite precision. In this framework, macroscopic irreversibility emerges from the overwhelming statistical likelihood that a system will evolve toward states of higher entropy, even though the underlying equations are time-symmetric.

The Ehrenfest urn model [26] provides an elementary probabilistic illustration suggesting that entropy, viewed microphysically, might even decrease. Consider two urns initially containing an uneven distribution of balls, with most in the first urn. Assume a constant probability per time step for any ball to transfer to the other urn. As the system evolves, it will most likely approach an equilibrium state with roughly equal numbers of balls (a 50:50 ratio) in both urns, corresponding to maximum entropy in this analogy. However, even from this high-probability equilibrium state, fluctuations are possible. Poincaré's recurrence theorem implies that after an ‘enormously long time,’ it is not only possible but inevitable that the system will return to highly improbable states, such as having all balls collected in a single urn. Therefore, the eventual reappearance of these low-entropy configurations (‘outliers’ or *Buckel*[26]) cannot be ruled out; indeed, their absence over sufficiently long timescales would be extremely improbable. This guaranteed recurrence forms the basis of Zermelo's *Wiederkehreinwand* (recurrence objection) against monotonic entropy increase at maximal (microphysical) resolution. The simulation results depicted in Figure 1 demonstrate the system's tendency towards a state of higher entropy (equilibrium), punctuated by fluctuations that manifest as temporary, occasional decreases in entropy. Moreover, if the system is capable of universal computation, recurrence times for certain ‘computationally complex, resource-intensive’ states—such as those associated with the halting probability Omega mentioned earlier—can be expected to grow ‘beyond any specifiable size’, potentially faster than any recursive (computable) lower bound [27].

In statistical mechanics, a macroscopic state is essentially an equivalence class, where the equivalence relation is defined




FIG. 1. Evolution of the number of balls in Urn 1 (blue) and Urn 2 (dotted, orange) in the Ehrenfest Urn Model ($N=100$, 1000 steps), starting from a low-entropy state with Urn 1 filled and Urn 2 empty. The dashed line marks the equilibrium state ($N/2=50$). The simulation highlights the system’s relaxation towards equilibrium and the persistent fluctuations around it, illustrating the microscopic reversibility that underlies Zermelo’s recurrence objection.

by the condition that two microscopic states are considered equivalent if they share the same values for macroscopic variables, such as energy or volume. In the aforementioned example, macroscopic states with roughly equal numbers of balls (a 50:50 ratio) in both urns are much more likely than macroscopic states characterized by outliers with the same number of balls. Therefore, macroscopic systems tend to evolve toward entropy increase.

For example, in a gas, all possible molecular arrangements that result in the same pressure and temperature belong to the same macroscopic state. Consequently, microscopic states can be formally ‘bundled together’ or ‘grouped’ based on macroscopic equivalence: If they cannot be distinguished through operational means at the macroscopic level, they are defined as equivalent. The corresponding binary equivalence relation, applied to microstates, naturally satisfies the properties of reflexivity, symmetry, and transitivity.

In summary, while the mathematical description of a system in terms of infinite precision microstates leads to reversible trajectories, the physical impossibility of achieving such precision guarantees that real systems display irreversible behavior. The irreversible macroscopic laws of thermodynamics are thus understood as emergent, effective descriptions that arise from practical limitations on precision and the statistical averaging over an enormous number of microstates. The physical means define an equivalence relation on microphysical states. The corresponding equivalence classes can be identified with macroscopic states.

IV. FORMALIZATION OF FAPPNESS BY EQUIVALENCE RELATIONS

In an early critique of sectorization-type arguments [28, 29] reviewed in Section IID, Bell argued [30] that unlimited or even actually infinite means are physically unattainable. He later introduced the related concept of For-All-Practical-Purposes (FAPP) [31], which replaces transfinite means with finite, physically operational ones.

The concept of FAPP indistinguishability can be rigorously formalized using equivalence relations. In different physical contexts, these equivalence relations partition microscopic configurations into equivalence classes, grouping together states that are operationally indistinguishable at a higher level of description. The three primary instantiations of such equivalence classes, as discussed in previous sections, are:

A. Classical Analysis

In classical mechanics and dynamical systems, coarse-graining leads to an effective partitioning of phase space into equivalence classes. Two microstates belong to the same class if they yield identical macroscopic observables within a given resolution limit. This follows naturally from measurement constraints and computational limitations in practical analysis.

B. Sectorization in Quantum Mechanics

In quantum theory, the emergence of classical-like behavior is often described using *superselection sectors* or *decoherence-induced equivalence classes*. Here, quantum states that differ only by superpositions within a decohered basis (due to environmental interactions) become practically indistinguishable. Such states effectively belong to the same equivalence class, as they do not interfere and cannot be resolved through macroscopic measurements.

C. Macrostates in Statistical Physics

In statistical mechanics, a macroscopic state corresponds to an equivalence class of microstates that share the same macroscopic variables, such as energy, volume, or magnetization. Since individual microstates fluctuate rapidly and are inaccessible in practice, all configurations that yield the same macroscopic properties are grouped together, forming a thermodynamic macrostate.

In all three cases, the corresponding equivalence relation on microstates satisfies reflexivity, symmetry, and transitivity, ensuring a well-defined partitioning of state space. This formalization captures the essence of FAPP reasoning, where practical indistinguishability justifies the use of equivalence classes in physical descriptions.

V. CONCLUSION

This paper explored the role of infinity in bridging microscopic and macroscopic descriptions in physics, focusing on the emergence of irreversibility from reversible dynamics. We examined how infinite processes are essential in mathematical constructions, such as the transition from rational to real numbers, and how they manifest in physical theories, from statistical mechanics to quantum measurement.

In classical analysis, infinite precision is a theoretical idealization that is unattainable in practice. The necessity of coarse-graining and finite resolution in measurements leads naturally to the formation of equivalence classes that group together states indistinguishable for all practical purposes (FAPP). This provides a foundation for understanding macro-

scopic irreversibility despite the underlying time-reversible microscopic laws.

In quantum mechanics, infinite tensor products and sectorization offer a framework for understanding the transition from unitary evolution to apparent wavefunction collapse. Von Neumann's insights [18] emphasize that the full set of product states in an infinite tensor product is uncountably infinite, with a cardinality of 2^{\aleph_0} . The resulting space is nonseparable. This sharply contrasts with any countable subset, which has a cardinality of \aleph_0 ; here, the distinction reflects the difference between nonseparability and separability. Since unitary operators preserve the inner product structure—and, consequently, the cardinality of any orthonormal basis—no unitary transformation can map a countable subset onto the full uncountable set. Therefore, under constraints such as (finite or infinite) denumerable group actions, unitary equivalence fails in the limit of infinite tensor products. The partitioning of Hilbert space into equivalence classes through decoherence-induced superselection rules highlights how quantum-to-classical transitions can emerge from infinite degrees of freedom.

In statistical mechanics, macrostates are equivalence classes of microstates that share the same macroscopic observables, such as energy or volume. The practical impossibility of resolving individual microstates supports the statistical interpretation of thermodynamic irreversibility.

By formalizing Bell's FAPP approach using equivalence relations, we provided a unifying perspective on how operational indistinguishability underlies emergent macroscopic behavior. Across classical analysis, quantum mechanics, and statistical physics, equivalence classes play a crucial role in describing physical reality at different scales. This perspective underscores the foundational role of infinity in physics and its implications for the nature of measurement, irreversibility, and emergent phenomena.

ACKNOWLEDGMENTS

The author gratefully acknowledges discussions with Nisson S. Yanofsky. I am grateful to an anonymous referee for drawing my attention to surreal numbers in this context. This research was funded in whole or in part by the Austrian Science Fund (FWF) [Grant DOI:10.55776/I4579].

[1] J. H. Conway, *On Numbers and Games*, 2nd ed. (A K Peters, Ltd., Natick, MA, USA, 2001).

[2] D. E. Knuth, *Surreal Numbers* (Addison-Wesley, Reading, MA, USA, 1974).

[3] N. S. Yanofsky, A universal approach to self-referential paradoxes, incompleteness and fixed points, *Bulletin of Symbolic Logic* **9**, 362 (2003), arXiv:math/0305282.

[4] P. W. Bridgman, A physicist's second reaction to Mengenlehre, *Scripta Mathematica* **2**, 101 (1934).

[5] K. Svozil, Set theory and physics, *Foundations of Physics* **25**, 1541 (1995).

[6] N. S. Yanofsky, private communication (2025), dated March 6, 2025.

[7] E. Specker, Nicht konstruktiv beweisbare Sätze der Analysis, *The Journal of Symbolic Logic* **14**, 145 (1949).

[8] G. Kreisel, A notion of mechanistic theory, *Synthese* **29**, 11 (1974).

[9] P. Ehrenfest, Graphische Veranschaulichung des einfachsten Falles von ungleichförmiger Reihenkonvergenz, *Mathematisch-Naturwissenschaftliche Blätter* **6**, 1 (1909).

[10] G. J. Chaitin, A theory of program size formally identical to information theory, *Journal of the Association of Computing*

Machinery (JACM) **22**, 329 (1975).

[11] C. S. Calude and M. J. Dinneen, Exact approximations of omega numbers, International Journal of Bifurcation and Chaos **17**, 1937 (2007), CDMTCS report series 293.

[12] G. J. Chaitin, Computing the busy beaver function, in *Open Problems in Communication and Computation*, edited by T. M. Cover and B. Gopinath (Springer, New York, 1987) p. 108.

[13] P. Martin-Löf, The definition of random sequences, Information and Control **9**, 602 (1966).

[14] C. S. Calude, *Information and Randomness—An Algorithmic Perspective*, 2nd ed. (Springer, Berlin, 2002).

[15] J. von Neumann, *Mathematische Grundlagen der Quantenmechanik*, 2nd ed. (Springer, Berlin, Heidelberg, 1932, 1996) English translation in [16].

[16] J. von Neumann, *Mathematical Foundations of Quantum Mechanics*, Princeton Landmarks in Mathematics and Physics (Princeton University Press, Princeton, NJ, USA, 2018) translated by Robert T. Beyer, edited by Nicholas A. Wheeler.

[17] H. Everett III, ‘Relative State’ formulation of quantum mechanics, Reviews of Modern Physics **29**, 454 (1957).

[18] J. von Neumann, On infinite direct products, Compositio Mathematica **6**, 1 (1939), reprinted in *John von Neumann, Collected Works, Vol. III*, A. H. Taub, editor, Pergamon Press, New York, 1961, nr. 6, p. 323–399.

[19] P. Grangier, Completing the quantum formalism in a contextually objective framework, Foundations of Physics **51**, 10.1007/s10701-021-00424-1 (2021), arXiv:2003.03121.

[20] M. Van Den Bossche and P. Grangier, Contextual unification of classical and quantum physics, Foundations of Physics **53**, 10.1007/s10701-023-00678-x (2023), arXiv:2209.01463.

[21] M. Van Den Bossche and P. Grangier, Postulating the unicity of the macroscopic physical world, Entropy **25**, 1600 (2023).

[22] M. Van Den Bossche and P. Grangier, Revisiting quantum contextuality in an algebraic framework, Journal of Physics. Conference Series **2533**, 012008 (2023), arXiv:2304.07757.

[23] O. Darrigol, Boltzmann’s reply to the Loschmidt paradox: A commented translation, The European Physical Journal H **46**, 10.1140/epjh/s13129-021-00029-2 (2021).

[24] J. C. Maxwell, Report on a paper by Prof. Osborne Reynolds ‘on certain dimensional properties of matter in the gaseous state’ (1897), dated 28 March, 1879, Royal Society of London, Archives, no. 188, Reference number: RR/8/188, reprinted in [25].

[25] E. Garber, S. G. Brush, and C. W. F. Everitt, *Maxwell on Heat and Statistical Mechanics: On “Avoiding All Personal Enquiries” of Molecules* (Lehigh University Press and Associated University Press, Bethlehem and London, 1995).

[26] P. Ehrenfest and T. Ehrenfest, Über eine Aufgabe aus der Wahrscheinlichkeitsrechnung, die mit der kinetischen Deutung der Entropievermehrung zusammenhängt, Mathematisch-Naturwissenschaftliche Blätter **3**, 1 (1906).

[27] K. Svozil, *Randomness & Undecidability in Physics* (World Scientific, Singapore, 1993).

[28] K. Hepp, Quantum theory of measurement and macroscopic observables, Helvetica Physica Acta **45**, 237 (1972).

[29] J. Bub, The measurement problem from the perspective of an information-theoretic interpretation of quantum mechanics, Entropy **17**, 7374 (2015).

[30] J. S. Bell, On wave packet reduction in the Coleman-Hepp, Helvetica Physica Acta **48**, 93 (1975).

[31] J. S. Bell, Against ‘measurement’, Physics World **3**, 33 (1990).