

# A Quantum Mechanical Look at Time Travel and Free Will

Daniel Greenberger and Karl Svozil

## Abstract

Consequences of the basic and most evident consistency requirement – that measured events cannot happen and not happen at the same time – are reviewed. Particular emphasis is given to event forecast and event control. As a consequence, particular, very general bounds on the forecast and control of events within the known laws of physics result. These bounds are of a global, statistical nature and need not affect singular events or groups of events. We also present a quantum mechanical model of time travel and discuss chronology protection schemes. Such models impose restrictions upon particular capacities of event control.

## 1 Classical Part

### 1.1 Principle of Self-Consistency

An irreducible, atomic physical phenomenon manifests itself as a click of some detector. Either there is a click or there is no click. This yes-no scheme is experimental physics in a nutshell (at least according to a theoretician). From this kind of elementary observation, all of our physical evidence is accumulated.

Such irreversibly observed events (whatever the relevance or meaning of those terms is (Wigner 1961, Wheeler 1983, Greenberger and YaSin 1989, Herzog et al. 1995)) are subject to the primary condition of *consistency* or *self-consistency*. *Any particular irreversibly observed event either happens or does not happen, but it cannot both happen and not happen.*

Indeed, so trivial seems the requirement of consistency that Hilbert polemicized against “another author” with the following words (Hilbert 1926), “... for me, the opinion that the [physical] facts and events themselves can be contradictory is a good example of thoughtlessness.”

Just as in mathematics, inconsistency, i.e. the coexistence of truth and falsity of propositions, is a fatal property of any physical theory. Nevertheless, in a particular very precise sense, quantum mechanics incorporates inconsistencies in a very subtle way, which assures overall consistency. For instance, a particle wave function or quantum state is said to “pass” a double slit through both slits at once, which is classically impossible. (Such considerations may, however, be considered as mere trickery, quantum talk devoid of any operational meaning.) Yet, neither particle wave functions nor quantum states can be directly associated with any sort of irreversible observed event of physical reality. We shall come back to a particular quantum case in the second part of this investigation.

And just as in mathematics and in formal logic, it can be argued that overly strong capacities of intrinsic event forecast and intrinsic event control render the system overall inconsistent. This fact may indeed be considered as one decisive feature in finite deterministic (“algorithmic”) models (Svozil 1993). It manifests itself already in the early stages of Cantorian set theory: any claim that it is possible to enumerate the real numbers leads, via the diagonalization method, to an outright contradiction. The only consistent alternative is the acceptance that no such capacity of enumeration exists. Gödel’s incompleteness theorem (Gödel 1931) states that any formal system rich enough to include arithmetic and elementary logic could not be both consistent and complete. Turing’s theorem on the recursive unsolvability of the halting problem (Turing 1936/1937), as well as Chaitin’s  $\Omega$  numbers (Chaitin 1992) are formalizations of related limitations in formal logics, computer science and mathematics.

In what follows we will proceed along very similar lines. We will first argue that any capacity of total forecast or event control – even in a totally deterministic environment – contradicts the (idealistic) idea that decisions between alternatives are possible; or, stated differently, that there is free will. Then we shall proceed with possibilities of forecast and event control which are consistent with both free will and the known laws of physics.

Evidently, some form of forecast and event control is possible – indeed, this is one of the main achievements of contemporary natural science, and we make everyday use of it, e.g. by switching on the light. These capacities of forecast and event control are characterized by a high degree of reproducibility, which does not depend on single events.

We will concentrate on very general bounds for these capacities, which follow from the requirement of consistency and do not depend on any particular physical model. They are valid for all conceivable forms of physical theories – classical, quantum and forthcoming alike.

## 1.2 Strong Forecasting

Let us first consider forecasting the future. Even if physical phenomena occur deterministically and can be accounted for (“computed”) on a higher level of abstraction, from within the system such a complete description may not be of much practical, operational use (Toffoli 1978, Svozil 1996).

Indeed, suppose that free will exists. Suppose further that an agent could predict *all* future events, without exceptions. We shall call this the *strong form of forecasting*. In this case, the agent could freely decide to act in such a way as to invalidate any prediction. Hence, in order to avoid inconsistencies and paradoxes, either free will has to be abandoned, or it has to be accepted that complete prediction is impossible.<sup>1</sup>

Another possibility would be to consider strong forms of forecasting which are, however, not utilized to alter the system. Effectively, this results in the abandonment of free will, amounting to an extrinsic, detached viewpoint. After all, what is knowledge and what is it good for if it cannot be applied and utilized?

Recent advances in the foundations of quantum (information) theory have shown that, due to complementarity and the impossibility to clone generic states, single events may have important meanings to some observers, although they make no sense at all to other observers. One example for this is quantum cryptography. Many of these events are stochastic and are postulated to satisfy all conceivable statistical laws (correlations are nonclassical, though). In such frameworks, high degrees of reproducibility cannot be guaranteed, although single events may carry valuable information, which can even be distilled and purified.

## 1.3 Strong Event Control

A very similar argument holds for event control and the production of “miracles” (Frank 1932). Suppose that free will exists. Suppose further that an agent could entirely control the future. We will call this the *strong form of event control*. Then this agent could freely decide to invalidate the laws of physics. In order to avoid a paradox, either free will or some physical laws would have to be abandoned, or it has to be accepted that complete event control is impossible.

---

<sup>1</sup>This argument is of an ancient type (Anderson 1970). As has already been mentioned, it has been formalized recently in set theory, formal logic and recursive function theory, where it is called the “diagonalization method.”

## 1.4 Weak Forecasting and Event Control

Already from what has been said, it is reasonable to assume that *forecast and event control should be possible only if these capacities cannot be associated with any paradox or contradiction.*

Thus the requirement for consistency of the phenomena seems to impose rather stringent conditions on forecasting and event control. Similar ideas have already been discussed in the context of time paradoxes in relativity theory (cf. Friedman et al. 1990 and Nahin 1998, p. 272: “the only solutions to the laws of physics that can occur locally ... are those which are globally self-consistent”).

There is, however, a possibility that the forecast and control of future events *is* conceivable for *singular* events within the statistical bounds. Such occurrences may be “singular miracles” which are well accounted for within the known laws of physics. They will be called *weak forms of forecasting and event control*. In order to obey overall consistency, such a framework should not be extendible to any forms of strong forecast or event control, because, as has been argued before, this could either violate global consistency criteria or would make necessary a revision of the known laws of physics.

The relevant laws of statistics (e.g. all recursively enumerable laws) impose rather lax constraints, especially on finite sequences, and do not exclude local, singular, improbable events. For example, a binary sequence such as 11111111111111111111111111111111 is just as probable as the sequences 11100101110101000111000011010101 and 01010101010101010101010101010101, and its occurrence in a test is equally likely, although the “meaning” an observer could ascribe to it is rather different. These sequences may be embedded in and be part of much longer stochastic sequences. If short finite regular (or “meaningful”) sequences are embedded into long irregular (“meaningless”) ones, those sequences become statistically indistinguishable for all practical purposes from the previous sequences. Of course, the “meaning” of any such sequence may vary with different observers. Some of them may be able to decipher a sequence, others may not be able to do so.

It may seem evident that by definition any finite regularity in an otherwise stochastic environment should exclude the type of high reproducibility characteristic of the natural sciences. On the contrary: single “meaningful” events, which are hardly reproducible, might indicate a new category of phenomena dual to the usual “lawful” and highly predictable ones.<sup>2</sup>

---

<sup>2</sup>In this context compare the contribution by Primas (particularly sections 6–8) in this volume.

Just as it is perfectly all right to consider the statement "This statement is true" to be true, it may be perfectly reasonable to speculate that some events are forecasted and controlled within the domain of statistical laws. But in order to be within the statistical laws, any such method *need not be guaranteed* to work at all times.

To put it pointedly: it may be perfectly reasonable to become rich, say, by singular forecasts of the stock market, future values or the outcomes of horse races, but such an ability must necessarily be irreproducible, secretive, and not extendible; at least to such an extent that no guarantee for an overall strategy and regularity can be derived from it.

The associated weak forms of forecasting and event control are thus beyond any global statistical significance. Their importance and meaning seems to lie mainly on a "subjective" level of singular events. This comes close to what Jung imagined as the principle of "synchronicity" (Jung 1952), and is dual to the more reproducible forms one is usually accustomed to.

### 1.5 Against the Odds

Let us review a couple of experiments which suggest themselves in the context of weak forecast and event control. They are all based on the observation whether or not an agent is capable of correctly forecasting or controlling future events such as, say, the tossing of a fair coin.

In the first run of such an experiment, no consequence is derived from the agent's capacities despite the mere recording of the data. The second run of the experiment is like the first run, but the *meaning* of the forecasts or controlled events is different. The events are taken as outcomes, e.g., of gambling against other individuals (i) with or (ii) without similar capacities, or against (iii) an anonymous "mechanical" agent such as a casino or a stock exchange. (As a variant of this experiment, the partners or adversaries of the agent are informed about the agent's intentions.)

In the third run of the experiment, the experimenter attempts to counteract the agent's capacities. Let us assume the experimenter has total control over the event. If the agent predicts or attempts to bring about a particular future event, the experimenter causes the event not to happen and so on.

It might be interesting to record just how much the agent's capacities are changed by the setup. An expectation might be defined from a dichotomic observable

$$e(A, i) = \begin{cases} +1 & \text{correct guess} \\ -1 & \text{incorrect guess} \end{cases}$$

where  $A$  stands for agent  $A$  and  $i$  stands for the  $i$ th experiment. An expecta-

tion function can then be defined as usual by the average over  $N$  experiments; i.e.

$$E(A) = \frac{1}{N} \sum_{i=1}^N e(A, i).$$

From the first to the second type of experiment it should become more and more unlikely that the agent operates correctly, since his performance is leveled against other agents with more or less the same capacities. The third type of experiment should produce a total anticorrelation. Formally, this should result in a decrease of  $E$  when compared to the first round of experiment.

Another, rather subtle deviation from probabilistic laws may be observed if *correlated* events are considered. Just as in the case of quantum entanglement, it may happen that individual components of correlated systems behave totally at random and exhibit more disorder than the system as a whole (Nielsen and Kempe 2001).

If once again one assumes two dichotomic observables  $e(A, i)$  and  $e(B, i)$  of a correlated subsystem, then the correlation function

$$C(A, B) = \frac{1}{N} \sum_{i=1}^N e(A, i) e(B, i)$$

and the associated probabilities may give rise to violations of the Boole-Bell inequalities – Boole’s *conditions of possible (classical) experience* (Boole 1862, Hailperin 1976, Pitowsky 1989, 1994) – and may even exceed (Krenn and Svozil 1998) the Tsirelson bounds (Cirel’son 1980, Tsirel’son 1987, Cirel’son 1993) for *conditions of possible (quantum) experience*. There, the agent should concentrate on influencing the *coincidences* of the event rather than the single individual events. In such a case, the *individual* observables may behave perfectly random, while the associated *correlations* might be nonclassical and even stronger-than-quantum, and might give rise to highly nonlocal phenomena. As long as the individual events cannot be controlled, this does not need to violate Einstein causality. (But, even then, consistent scenarios remain (Svozil 2000)).

In summary, it can be stated that, although total forecasting and event control are incompatible with free will, more subtle forms of these capacities remain conceivable even beyond the present laws of physics; at least as long as their effects upon the “fabric of phenomena” are consistent. These capacities are characterized by singular events and not by statistically reproducible patterns, which are often encountered under the known laws of

physics. Whether or not such capacities exist remains an open question. Nevertheless, despite the elusiveness of the phenomenology involved, it does not appear unreasonable that the hypothesis could be operationalized, tested and even put to use in particular contexts.

## 2 Quantum Part

### 2.1 Quantum Information

By coherent superposition, quantum theory manages to implement two classically inconsistent bits of information by one quantum bit. For example, consider the states  $|+\rangle$  and  $|-\rangle$  associated with the proposition that the spin of an electron in a particular direction is “up” or “down,” respectively. The coherent superposition of these two states  $(|+\rangle + |-\rangle)/\sqrt{2}$  is a 50:50 mixture of these two classically distinct possibilities and at the same time is a perfect quantum state.

Based upon this feature, we speculate that we may be able to solve some tasks which are classically intractable or even inconsistent by superposing quantum states in a self-consistent manner. In particular, we could speculate that diagonalization tasks using *not*-gates may become feasible, although the capacities of agents within such semi-closed time loops may be limited by requirements of (self-)consistency, which translate into bounds due to unitary quantum time evolution. These quantum consistency requirements, however, may be less restrictive than in the classical case (Svozil 1995a,b).

### 2.2 Mach-Zehnder Interferometer with Feedback Loop

In what follows we shall consider a Mach-Zehnder interferometer as drawn in Fig. 1 with two input and two output ports (Greenberger et al. 1993). The novel feature of this device is a feedback loop from the future of one output port into the past of an input port. Thereby we leave open the question of such a feedback loop into the past and how it can (if ever) be realized. Indeed, if one dislikes the idea of backwards-in-time communication, one may think of this feedback loop as a channel which, by synchronizing the beams, acts as if a beam from the future enters the input port, while this beam actually was emitted in the past from the output port.

If one merely introduced feedback as in classical electrical engineering, this would defy unitarity, as two input channels would be going into one forward channel, which could not be uniquely reversed. So one needs a feedback coupling that resembles a beam-splitter, as in Fig. 1. The operator




Figure 1: Mach-Zehnder device with backwards-in-time output  $\psi_4(t_2)$  which passes  $M$  and serves as input  $\psi_4(t_1)$ .

$M$  generates the effects of the feedback in time. These “beam-splitters” are figurative. Their role is to couple the two incoming channels to two outgoing channels. The operator  $G_1$  represents the ordinary time development in the absence of time feedback. The operator  $G_2$  represents an alternate possible time evolution that can take place and compete with  $G_1$  because there is feedback. We want to find ..... in the presence of the feedback in time that is generated by the operator  $M$ . At the beam splitters, the forward amplitude is  $\alpha$ , while the reflected amplitude is  $i\beta$ . The beam splitters are shown in Fig. 2. They perform the unitary transformation:

$$\begin{aligned} |a\rangle &= \alpha|d\rangle + i\beta|c\rangle \\ |b\rangle &= \alpha|c\rangle + i\beta|d\rangle \end{aligned} \quad (1)$$

Here we assume for simplicity that  $\alpha$  and  $\beta$  are real. We can invert this to obtain:

$$\begin{aligned} |d\rangle &= \alpha|a\rangle - i\beta|b\rangle \\ |c\rangle &= \alpha|b\rangle - i\beta|a\rangle \end{aligned} \quad (2)$$

The overall governing equations can be read from Fig. 2. At time  $t_2$  the second beam-splitter determines  $\psi_3(t_2)$  and  $\psi_4(t_2)$ . We have

$$\psi_3(t_2) \equiv \psi'_3 = \alpha\psi_1(t_2) - i\beta\psi_2(t_2) = \alpha\psi'_1 - i\beta\psi'_2, \quad (3)$$






Figure 2: Reflection and transmission through a mirror with reflection coefficient  $\beta$  and transmission coefficient  $\alpha$ .

where the prime indicates the time  $t_2$  in the argument, and no prime indicates the time  $t_1$ . The wave functions  $\psi_1$  and  $\psi_2$  are determined at time  $t_2$  by

$$\psi_1(t_2) \equiv \psi'_1 = G_1\psi_1(t_1) = G_1\psi_1, \quad (4)$$

$$\psi_2(t_2) \equiv \psi'_2 = G_2\psi_2(t_1) = G_2\psi_2, \quad (5)$$

so that from eq. (3),

$$\psi'_3 = \alpha G_1\psi_1 - i\beta G_2\psi_2, \quad (6)$$

and equivalently:

$$\psi'_4 = \alpha G_2\psi_2 - i\beta G_1\psi_1 \quad (7)$$

The propagator  $M$  is what produces the feedback in time, propagating from  $t_2$  back to  $t_1$ , so that  $\psi_4(t_1) = M\psi_4(t_2)$ , or

$$\psi_4 = M\psi'_4. \quad (8)$$

At the beamsplitter at  $t_1$ , we have:

$$\psi_1 = \alpha\psi - i\beta\psi_4, \quad (9)$$

$$\psi_2 = \alpha\psi_4 - i\beta\psi. \quad (10)$$

### 2.3 The Solution

First, we want to eliminate the  $\psi_4$  in eqs. (9) and (10), to get equations for  $\psi_1$  and  $\psi_2$ . Then from eq. (6) we can obtain  $\psi'_3$ . From eqs. (7) and (8), we have:

$$\psi_4 = M\psi'_4 = \alpha MG_2\psi_2 - i\beta MG_1\psi_1 \quad (11)$$

We plug this into eqs. (9) and (10),

$$\psi_1 = \alpha\psi - i\beta(\alpha MG_2\psi_2 - i\beta MG_1\psi_1), \quad (12)$$

$$\psi_2 = \alpha(\alpha MG_2\psi_2 - i\beta MG_1\psi_1) - i\beta\psi, \quad (13)$$

and rewrite these as:

$$\psi_1 = (1 + \beta^2 MG_1)^{-1}(-i\alpha\beta MG_2)\psi_2 + \alpha(1 + \beta^2 MG_1)^{-1}\psi, \quad (14)$$

$$\psi_2 = (1 - \alpha^2 MG_2)^{-1}(-i\alpha\beta MG_1)\psi_1 - i\beta(1 - \alpha^2 MG_2)^{-1}\psi. \quad (15)$$

These two simultaneous equations must be solved to find  $\psi_1$  and  $\psi_2$  as functions of  $\psi$ . To solve for  $\psi_1$ , substitute eq. (15) into (14) such that

$$\begin{aligned} \psi_1 = & (1 + \beta^2 MG_1)^{-1}(-i\alpha\beta MG_2)[(1 - \alpha^2 MG_2)^{-1}(-i\alpha\beta MG_1)\psi_1 \\ & - i\beta(1 - \alpha^2 MG_2)^{-1}\psi] + \alpha(1 + \beta^2 MG_1)^{-1}\psi \end{aligned} \quad (16)$$

or:

$$\begin{aligned} [1 + \alpha^2\beta^2(1 + \beta^2 MG_1)^{-1}(MG_2)(1 - \alpha^2 MG_2)^{-1}(MG_1)]\psi_1 \\ = (1 + \beta^2 MG_1)^{-1}[-\alpha\beta^2 MG_2(1 - \alpha^2 MG_2)^{-1} + \alpha]\psi \end{aligned} \quad (17)$$

If we rewrite this as

$$[X]\psi_1 = (Y)^{-1}[Z]\psi, \quad (18)$$

we can simplify the equation as:

$$\begin{aligned} XY &= 1 + \beta^2 MG_1 + \alpha^2\beta^2 MG_2(1 - \alpha^2 MG_2)^{-1}MG_1 \\ &= 1 + \beta^2[1 + (1 - \alpha^2 MG_2)^{-1}\alpha^2 MG_2]MG_1 \\ &= 1 + \beta^2(1 - \alpha^2 MG_2)^{-1}MG_1, \end{aligned} \quad (19)$$

and

$$\begin{aligned} Z &= \alpha(1 - \alpha^2 MG_2)^{-1}(1 - \alpha^2 MG_2 - \beta^2 MG_2) \\ &= \alpha(1 - \alpha^2 MG_2)^{-1}(1 - MG_2). \end{aligned} \quad (20)$$

Thus,

$$\psi_1 = \alpha[1 + \beta^2(1 - \alpha^2 MG_2)^{-1}MG_1]^{-1}(1 - \alpha^2 MG_2)^{-1}(1 - MG_2)\psi. \quad (21)$$

Then, using the identity  $A^{-1}B^{-1} = (BA)^{-1}$ , we finally obtain

$$\psi_1 = \alpha(1 - \alpha^2 MG_2 + \beta^2 MG_1)^{-1}(1 - MG_2)\psi. \quad (22)$$

We can solve for  $\psi_2$  similarly, by substituting eq. (14) into (15):

$$\psi_2 = -i\beta(1 - \alpha^2 MG_2 + \beta^2 MG_1)^{-1}(1 + MG_1)\psi. \quad (23)$$

Notice that in the denominator terms in eqs. (22) and (23),  $\alpha$  and  $\beta$  have reversed the role of the operators they apply to. We can finally use eq. (6) to solve for  $\psi'_3 = \psi_3(t_2)$ :

$$\psi_3(t_2) = [\alpha^2 G_1 D(1 - MG_2) - \beta^2 G_2 D(1 + MG_1)]\psi(t_1), \quad (24)$$

where  $D = (1 + \beta^2 MG_1 - \alpha^2 MG_2)^{-1}$ .

## 2.4 Important Special Cases

(i) For commuting  $M$ ,  $G_1$  and  $G_2$ ,  $D = \beta^2(1 + MG_1) + \alpha^2(1 - MG_2)$ , and

$$\psi'_3 = \frac{\alpha^2 G_1 - \beta^2 G_2 - MG_1 G_2}{1 + \beta^2 MG_1 - \alpha^2 MG_2} \psi(t_1). \quad (25)$$

(ii) For  $\alpha = 1$ ,  $\beta = 0$ , there is no feedback. Here

$$\psi'_3 = G_1(1 - MG_2)^{-1}(1 - MG_2)\psi = G_1\psi. \quad (26)$$

(iii) For  $\beta = 1$ ,  $\alpha = 0$ , there is only feedback. Here

$$\psi'_3 = -G_2(1 + MG_1)^{-1}(1 + MG_1)\psi = -G_2\psi. \quad (27)$$

(iv)  $G_1 = G_2 \equiv G$  :

$$\psi'_3 = G[1 + (\beta^2 - \alpha^2)MG]^{-1}(\alpha^2 - \beta^2 - MG)\psi. \quad (28)$$

(iv') If also  $\alpha^2 = \beta^2 = \frac{1}{2}$ , then

$$\psi'_3 = -GMG\psi. \quad (29)$$

(v) If  $\beta \ll 1$ , which is expected to be the usual case, then the solution only depends on  $\beta^2 = \gamma$ . Also,  $\alpha^2 = 1 - \beta^2 = 1 - \gamma$ . Then, to lowest order in  $\gamma$ , the denominator  $D$  in eq. (24) becomes

$$\begin{aligned} D &= [1 + \gamma MG_1 - (1 - \gamma)MG_2]^{-1} \\ &= (1 - MG_2)^{-1} - \gamma(1 - MG_2)^{-1}(MG_1 + MG_2)(1 - MG_2)^{-1} \end{aligned} \quad (30)$$

so that

$$\begin{aligned} \psi'_3 &= \{(1 - \gamma)G_1[1 - \gamma(1 - MG_2)^{-1}(MG_1 + MG_2)]\} \psi \\ &\quad - \{\gamma G_2(1 - MG_2)^{-1}(1 + MG_1)\} \psi \\ &= [G_1 - \gamma(G_1 + G_2)(1 - MG_2)(1 + MG_1)] \psi. \end{aligned} \quad (31)$$

(vi) The case that corresponds to the classical paradox that an agent shoots his father before he has met the agent's mother, so that the agent can never be born, has an interesting quantum-mechanical resolution. This is the case  $G_1 = 0$ , where there is a perfect absorber in the beam so that the system would never get to evolve to time  $t_2$ . But quantum mechanically, there is another path along  $G_2$ , at which the agent does not shoot his father, that has a probability  $\beta$  without feedback. The solution in this case is

$$\psi'_3 = -\beta^2 G_2(1 - \alpha^2 MG_2)^{-1} \psi. \quad (32)$$

We assume for simplicity that  $G_2$  is the standard time evolution operator

$$G_2 = e^{-iE(t_2-t_1)/\hbar}, \quad (33)$$

and  $M$  is the simplest backwards-in-time evolution operator

$$M = e^{-iE(t_1-t_2)/\hbar+i\varphi}, \quad (34)$$

where we have also allowed for an extra phase shift. Then

$$\begin{aligned} \psi'_3 &= -\beta e^{-iE(t_2-t_1)/\hbar} [1 - \alpha^2 e^{i\varphi}]^{-1} \psi, \\ |\psi'_3|^2 &= \frac{\beta^4}{(1 - \alpha^2 e^{i\varphi})(1 - \alpha^2 e^{-i\varphi})} |\psi|^2 \\ &= \frac{1}{1 + 4(\alpha^2/\beta^2) \sin^2(\varphi/2)} |\psi|^2. \end{aligned} \quad (35)$$

Note that for  $\varphi = 0$ ,  $\psi'_3 = -e^{-iE\Delta t/\hbar} \psi$  for *any* value of  $\beta$ . This means that no matter how small the probability that the agent ever reached here in the first place, the fact that he *is* here ( $\alpha \neq 1$ ) guarantees that, even though he is certain to have shot his father if he had met him ( $G_1 = 0$ ), the agent will not have met him! The agent will have taken the other path with 100% certainty.

How can we understand this result? In our model, with  $\varphi = 0$ , we have  $G_1 = 0$ , and  $MG_2 = 1$ . Also, we will assume that  $\beta \ll 1$ , even though this is not necessary. The various amplitudes are

$$\begin{aligned} |\psi_1| &= 0, & |\psi_2/\psi| &= 1/\beta, \\ |\psi_4/\psi| &= \alpha/\beta, & |\psi'_3/\psi| &= 1. \end{aligned} \quad (36)$$

So we see that the two paths of the beam-splitter at  $t_1$  leading to the path  $\psi_1$  cancel out. But of the beam  $\psi$ ,  $\alpha$  passes through, while of the beam  $\psi_4$ , only  $\beta$  leaks through. So the beam  $\psi_4$  must have a very large amplitude, which it does, as we can see from eqs. (36). In fact, it has a much larger amplitude than the original beam. Similarly, in order that  $|\psi'_3| = |\psi|$ ,  $\psi_2$  must have a very large amplitude. Thus we see that there is a large current flowing around the system, between  $\psi_2$  and  $\psi_4$ . But does this not violate unitarity? The answer is that if they were both running forward in time, it would. But one of these currents is running forward in time, while the other runs backward in time, and so they do not in this case violate unitarity. This is how our solution is possible.

So, according to our quantum model, if one could travel into the past, one would only see those alternatives consistent with the world one left. In

other words, while one could see the past, one could not change it. No matter how unlikely the events are that could have led to one's present circumstances, once they have actually occurred, they cannot be changed. One's trip would set up resonances that are consistent with the future that has already unfolded.

This also has consequences for the paradoxes of free will. It shows that it is perfectly logical to assume that one has many choices and that one is free to take any one of them. Until a choice is taken, the future is not determined. However, once a choice is taken, it was inevitable. It could not have been otherwise. So, looking backwards, the world is deterministic. However, looking forwards, the future is probabilistic.

The model also has consequences concerning a many worlds interpretation of quantum theory. The world may appear to keep splitting so far as the future is concerned, however once a measurement is made, only those histories consistent with that measurement are possible. In other words, with time travel, other alternative worlds do not exist, as once a measurement has been made, they would be impossible to reach from the original one.

Another interesting point comes from examining eq. (35). For small angles  $\varphi$  we see that

$$|\psi'_3|^2 = \frac{1}{1 + 4\frac{\alpha^2}{\beta^4} \sin^2(\varphi/2)} |\psi|^2 \rightarrow \frac{1}{-1 + \frac{\alpha^2 \varphi^2}{\beta^4}} |\psi|^2, \tag{37}$$

so that the above result is strongly resonant, with a Lorentzian shape, and a width  $\Delta\varphi \approx \beta^2$ , since  $\alpha \approx 1$ . Thus less "deterministic" and fuzzier time-travelling might be possible.

(vii) Sustained case: if we require the input and output state to be identical, i.e.  $\psi_3(t_2) = \psi(t_1)$ , then we obtain a sustainment condition (for commuting  $M, G_1, G_2$ ) of

$$1 = G_1(\alpha^2 - \beta^2 M) + G_2(\alpha^2 M - \beta^2) - MG_1 G_2. \tag{38}$$

Another case is  $G_1 = G_2 = 1$ , a phase shift in  $M = e^{i\varphi}$ , and  $\alpha = \beta = 1/\sqrt{2}$ , for which we obtain  $|\psi'_3| = |\psi|$ . For  $\beta = \sqrt{1 - \alpha^2} = 1/4$ ,

$$|\psi'_3| = \frac{112 - 113 \cos \varphi - 15i \sin \varphi}{54|1 - 7e^{i\varphi}/8|^2} |\psi|. \tag{39}$$

We summarize by stating that the structure of a quantum time travel through a Mach-Zehnder device is rich and unexpectedly elaborate. This suggests totally new szenarios for the possibility of free will and the capacities available to an agent acting in such a time loop.

## References

- Anderson A.R. (1970): Paul's epistle to Titus. In *The Paradox of the Liar*, ed. by R.L. Martin, Yale University Press, New Haven. The Bible contains a passage which refers to Epimenides, a Crete living in the capital city of Cnossus: "One of themselves, a prophet of their own, said, 'Cretans are always liars, evil beasts, lazy gluttons'." St. Paul, Epistle to Titus I (12–13).
- Boole G. (1862): On the theory of probabilities. *Philosophical Transactions of the Royal Society of London* **152**, 225–252.
- Chaitin G.J. (1992): *Information-Theoretic Incompleteness*, World Scientific, Singapore.
- Cirel'son B.S. (1980): Quantum generalizations of Bell's inequality. *Letters in Mathematical Physics* **4**, 93–100.
- Tsirel'son B.S. (= Cirel'son) (1987): Quantum analogues of the Bell inequalities. The case of two spatially separated domains. *Journal of Soviet Mathematics* **36**(4), ...–....
- Csirel'son B.S. (1993): Some results and problems on quantum Bell-type inequalities. *Hadronic Journal Supplement* **8**, 329–345.
- Davis M. (1965): *The Undecidable*, Raven Press, New York.
- Feferman S., Dawson J.W., Kleene S.C., Moore G.H., Solovay R.M., and van Heijenoort J., eds. (1986): *Kurt Gödel, Collected Works. Publications 1929–1936. Volume I*, Oxford University Press, Oxford.
- Frank P. (1932): *Das Kausalgesetz und seine Grenzen*, Springer, Vienna.
- Friedman J., Morris M.-S., Novikov I.-D., Echeverria F., Klinkhammer G., Thorne K.S., and Yurtsever U. (1990): Cauchy problem in spacetimes with closed timelike curves. *Physical Review D* **42**, 1915–1930.
- Gödel K. (1931): Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme. *Monatshefte für Mathematik und Physik* **38**, 173–198. English translation in Feferman et al. (1986) and in Davis (1965).
- Greenberger D.B. and YaSin A. (1989): "Haunted" measurements in quantum theory. *Foundation of Physics* **19**, 679–704.
- Greenberger D.B., Horne M., and Zeilinger A. (1993): Multiparticle interferometry and the superposition principle. *Physics Today* **46**, 22–29.
- Hailperin T. (1976): *Boole's Logic and Probability*, North-Holland, Amsterdam.

- Herzog T.J., Kwiat P.G., Weinfurter H., and Zeilinger A. (1995): Complementarity and the quantum eraser. *Physical Review Letters* **75**, 3034–3037.
- Hilbert D. (1926): Über das Unendliche. *Mathematische Annalen* **95**, 161–190.
- Jung C.G. (1952): Synchronizität als ein Prinzip akausaler Zusammenhänge. In *Naturerklärung und Psyche*, ed. by C.G. Jung and W. Pauli, Rascher, Zürich.
- Krenn G. and Svozil K. (1998): Stronger-than-quantum correlations. *Foundations of Physics* **28**, 971–984.
- Nahin P.J. (1998): *Time Travel*, second edition, Springer, New York.
- Nielsen M.A. and Kempe J. (2001): Separable states are more disordered globally than locally. *Physical Review Letters* **86**, 5184–5187.
- Pitowsky I. (1989): *Quantum Probability – Quantum Logic*, Springer, Berlin.
- Pitowsky I. (1994): George Boole’s ‘conditions of possible experience’ and the quantum puzzle. *Brit. J. Phil. Sci.* **45**, 95–125.
- Svozil K. (1993): *Randomness & Undecidability in Physics*, World Scientific, Singapore.
- Svozil K. (1995a): On the computational power of physical systems, undecidability, the consistency of phenomena and the practical uses of paradoxa. In *Fundamental Problems in Quantum Theory: A Conference Held in Honor of Professor John A. Wheeler. Annals of the New York Academy of Sciences* **755**, ed. by D. M. Greenberger and A. Zeilinger, Academy of Sciences, New York, pp. 834–841.
- Svozil K. (1995b): Consistent use of paradoxes in deriving constraints on the dynamics of physical systems and of no-go-theorems. *Foundations of Physics Letters* **8**, 523–535.
- Svozil K. (1996): Undecidability everywhere? In *Boundaries and Barriers. On the Limits to Scientific Knowledge*, Addison-Wesley, Reading, pp. 215–237.
- Svozil K. (2000): Relativizing relativity. *Foundations of Physics* **30**, 1001–1016.
- Toffoli T. (1978): The role of the observer in uniform systems. In *Applied General Systems Research*, ed. by G. Klir, Plenum Press, New York, pp. ...–

Turing A.M. (1936/1937): On computable numbers, with an application to the Entscheidungsproblem. *Proceedings of the London Mathematical Society, Series 2*, **42** and **43**, 230–265 and 544–546. Reprinted in Davis (1965).

Wheeler J.A. and Zurek W.H., eds. (1983): *Quantum Theory and Measurement*, Princeton University Press, Princeton.

Wheeler J.A. (1983): Law without law. In *Quantum Theory and Measurement*, ed. by J.A. Wheeler and W.H. Zurek, Princeton University Press, Princeton, pp. 182–213.

Wigner E.P. (1961): Remarks on the mind-body question. In *The Scientist Speculates*, ed. by I.J. Good Basic Books, New York, pp. 284–302. Reprinted in Wheeler and Zurek (1983) pp. 168–181.



**Between Chance and Choice**  
Interdisciplinary Perspectives  
on Determinism

Edited by  
Harald Atmanspacher and Robert Bishop

Imprint Academic  
Thorverton 2002

Copyright ©Imprint Academic, 2002

The moral rights of the authors have been asserted.

No part of any contribution may be reproduced in any form without permission, except for the quotation of brief passages in criticism and discussion.

Published in the UK by Imprint Academic  
PO Box 1, Thorverton EX5 5YX, UK

Published in the USA by Imprint Academic  
Philosophy Documentation Center  
PO Box 7147, Charlottesville, VA 22906-7147, USA

ISBN 0 907845 21 5

British Library Cataloguing in Publication Data  
A catalogue record for this book is available from the British Library.

Library of Congress Control Number: 2002102389