

Generalizing Tsirelson's Bound on Bell Inequalities Using a Min-Max Principle

Stefan Filipp*

Atominstytut der Österreichischen Universitäten, Stadionallee 2, A-1020 Vienna, Austria

Karl Svozil†

Institut für Theoretische Physik, University of Technology Vienna, Wiedner Hauptstraße 8-10/136, A-1040 Vienna, Austria

(Received 24 March 2004; published 23 September 2004)

Bounds on the norm of quantum operators associated with classical Bell-type inequalities can be derived from their maximal eigenvalues. This quantitative method enables detailed predictions of the maximal violations of Bell-type inequalities.

DOI: 10.1103/PhysRevLett.93.130407

PACS numbers: 03.65.Ud, 03.67.Mn

The violations of Bell-type inequalities represent a cornerstone of our present understanding of quantum probability theory [1]. Thereby, the usual procedure is as follows: First, the (in)equalities bounding the classical probabilities and expectations are derived systematically, e.g., by enumerating all conceivable classical possibilities and their associated two-valued measures. These form the extreme points which span the classical correlation polytopes [2–12], the faces of which are expressed by Bell-type inequalities which characterize the bounds of the classical probabilities and expectations—in Boole's term [13,14], the “conditions of possible experience.” (Generating functions is another method to find bounds on classical expectations [15,16].) The Bell-type inequalities contain sums of (joint) probabilities and expectations. In a second step, the classical probabilities and expectations in the Bell-type inequalities are substituted by quantum probabilities and expectations. The resulting operators violate the classical bounds. Until recently, little was known about the fine structure of the violations. Tsirelson (also written Cirel'son) published an absolute bound for the violation of a particular Bell-type inequality, the Clauser-Horne-Shimony-Holt (CHSH) inequality [2,3,17,18]. Cabello has investigated a violation of the CHSH inequality beyond the quantum mechanical bound by applying selection schemes to particles in a Greenberger-Horne-Zeilinger state [19,20]. Recently, detailed numerical [21] and analytical studies [22] stimulated experiments [23] to test the quantum bounds of certain Bell-type inequalities.

In what follows, a general method to compute quantum bounds on Bell-type inequalities is reviewed systematically. It makes use of the *min-max principle* for self-adjoint transformations (Ref. [24], Sec. 90 and Ref. [25], Sec. 75) stating that the operator norm is bounded by the minimal and maximal eigenvalues. These ideas are not entirely new and have been mentioned previously [15,21,22], yet to our knowledge no systematic investigation has been undertaken yet. It should also be kept in mind that this method *a priori* cannot produce quantum polytopes [21,26], but the quantum correspondents of

classical polytopes. Indeed, as we demonstrate explicitly, the resulting geometric forms are not convex. This, however, does not diminish the relevance of these quantum predictions to experiments testing the quantum violations of classical Bell-type inequalities.

As a starting point note that since $(A + B)^\dagger = A^\dagger + B^\dagger = (A + B)$ for arbitrary self-adjoint transformations A, B , the sum of self-adjoint transformations is again self-adjoint. That is, all self-adjoint transformations entering the quantum correspondent of any Bell-type inequality are again self-adjoint transformations. The sum does not preserve eigenvectors and eigenvalues; i.e., $A + B$ can have different eigenvectors and eigenvalues than A and B taken separately (i.e., A and B need not necessarily commute). The norm of the self-adjoint transformation resulting from summing the quantum counterparts of all the classical terms contributing to a particular Bell inequality obeys the min-max principle. Thus determining the maximal violations of classical Bell inequalities amounts to solving an eigenvalue problem. The associated eigenstates are the multipartite states which yield a maximum violation of the classical bounds under the given experimental (parameter) setup [27].

Let us demonstrate the method by considering two particles propagating in inverse directions, their polarization or spin being measured along two or more (m) distinct directions per particle perpendicular to their propagation directions. For these configurations, we enumerate analytical quantum bounds corresponding to the Clauser-Horne (CH) inequality, as well as of more general inequalities for $m > 2$ [10–12].

For $m = 2$, the CH inequalities restrict classical probabilities by $-1 \leq p_{13} + p_{14} + p_{23} - p_{24} - p_1 - p_3 \leq 0$, as well as permutations thereof. Here, p_1 and p_3 stand for the probabilities that the first particle is measured along the first direction and the second particle is measured along the third direction. p_{ij} stands for the joint probability to find the first particle along the direction i and the second particle along the direction j .

In order to evaluate the quantum counterpart of the CH inequalities, the classical probabilities have to be substi-

tuted by the quantum ones, i.e.,

$$\begin{aligned} p_1 &\rightarrow q_1(\theta) = \frac{1}{2}[\mathbb{1}_2 + \sigma(\theta)] \otimes \mathbb{1}_2, \\ p_3 &\rightarrow q_3(\theta) = \mathbb{1}_2 \otimes \frac{1}{2}[\mathbb{1}_2 + \sigma(\theta)], \\ p_{ij} &\rightarrow q_{ij}(\theta, \theta') = \frac{1}{2}[\mathbb{1}_2 + \sigma(\theta)] \otimes \frac{1}{2}[\mathbb{1}_2 + \sigma(\theta')], \end{aligned} \quad (1)$$

$$\sigma(\theta) = \begin{pmatrix} \cos\theta & \sin\theta \\ \sin\theta & -\cos\theta \end{pmatrix},$$

where θ is the relative measurement angle in the x - z plane, and the two particles propagate along the y axis. The quantum transformation associated with the CH inequality enumerated above is thus given by

with

$$\begin{aligned} O_{22}(\alpha, \beta, \gamma, \delta) &= q_{13}(\alpha, \gamma) + q_{14}(\alpha, \delta) + q_{23}(\beta, \gamma) - q_{24}(\beta, \delta) - q_1(\alpha) - q_3(\gamma) \\ &= \frac{1}{2}[\mathbb{1}_2 + \sigma(\alpha)] \otimes \frac{1}{2}[\mathbb{1}_2 + \sigma(\gamma)] + \frac{1}{2}[\mathbb{1}_2 + \sigma(\alpha)] \otimes \frac{1}{2}[\mathbb{1}_2 + \sigma(\delta)] + \frac{1}{2}[\mathbb{1}_2 + \sigma(\beta)] \\ &\quad \otimes \frac{1}{2}[\mathbb{1}_2 + \sigma(\gamma)] - \frac{1}{2}[\mathbb{1}_2 + \sigma(\beta)] \otimes \frac{1}{2}[\mathbb{1}_2 + \sigma(\delta)] - \frac{1}{2}[\mathbb{1}_2 + \sigma(\alpha)] \otimes \mathbb{1}_2 - \mathbb{1}_2 \otimes \frac{1}{2}[\mathbb{1}_2 + \sigma(\gamma)], \end{aligned} \quad (2)$$

where $\alpha, \beta, \gamma,$ and δ denote the measurement angles lying in the x - z plane: α and β for one particle, γ and δ for the other one. The eigenvalues of the self-adjoint transformation in (2) are

$$\lambda_{1,2,3,4}(\alpha, \beta, \gamma, \delta) = \frac{1}{2}[\pm\sqrt{1 \pm \sin(\alpha - \beta)\sin(\gamma - \delta)} - 1] \quad (3)$$

yielding the maximum bound $\|O_{22}\| = \max_{i=1,2,3,4} \lambda_i$. Note that for the particular choice of parameters $\alpha = 0, \beta = 2\theta, \gamma = \theta, \delta = 3\theta$ adopted in [21,22], one obtains $|O_{22}| = \frac{1}{2}\{[(3 - \cos 4\theta)/2]^{1/2} - 1\}$.

In the Bell basis $\{|\phi^+\rangle, |\psi^+\rangle, |\psi^-\rangle, |\phi^-\rangle\}$ with $|\psi^\pm\rangle = 1/\sqrt{2}(|01\rangle \pm |10\rangle)$ and $|\phi^\pm\rangle = 1/\sqrt{2}(|00\rangle \pm |11\rangle)$, the eigenvectors corresponding to the maximal violating eigenstates are

$$\begin{aligned} |\nu^\pm\rangle &= [F^\pm(\alpha, \beta, -\gamma, -\delta)|\psi^+\rangle + |\phi^-\rangle][1 + F^\pm(\alpha, \beta, -\gamma, -\delta)^2]^{-1/2}, \\ |\mu^\pm\rangle &= [F^\pm(\alpha, \beta, \gamma, \delta)|\phi^+\rangle + |\psi^-\rangle][1 + F^\pm(\alpha, \beta, \gamma, \delta)^2]^{-1/2}, \end{aligned} \quad (4)$$

with

$$F^\pm(\alpha, \beta, \gamma, \delta) = \pm 2\sqrt{1 - \sin(\alpha - \beta)\sin(\gamma - \delta)} \frac{\cos(\alpha - \delta) - \cos(\alpha - \gamma) - \cos(\beta - \gamma) - \cos(\beta - \delta)}{\sin(\alpha - \gamma) + \sin(\beta - \gamma) - \sin(\alpha - \delta) + \sin(\beta - \delta)}.$$

The states (4) are maximally entangled, corroborating the approach of Cabello [22] to utilize a set of maximally entangled states to reconstruct the quantum bound for the setting of the relative angles $\alpha = 0, \beta = 2\theta, \gamma = \theta,$ and $\delta = 3\theta$ [29]. From the particular form of the eigenstates, we conclude that the maximal violating eigenstates of the O_{22} operator are maximally entangled for general measurement angles lying in the x - z plane.

Generalizations for m measurements per particle are straightforward; for example, the extension to *three* measurement operators for each particle yields only one additional nonequivalent (with respect to symmetries) inequality [11,12] $I_{33} = p_{14} + p_{15} + p_{16} + p_{24} + p_{25} - p_{26} + p_{34} - p_{35} - p_1 - 2p_4 - p_5 \leq 0$ among the 684 inequalities [10] representing the faces of the associated classical correlation polytope. The associated operator for symmetric measurement directions is given by

$$\begin{aligned} O_{33}(0, \theta, 2\theta, 0, \theta, 2\theta) &= q_{14}(0, 0) + q_{15}(0, \theta) + q_{16}(0, 2\theta) + q_{24}(\theta, 0) + q_{25}(\theta, \theta) - q_{26}(\theta, 2\theta) + q_{34}(2\theta, \theta) - q_{35}(2\theta, \theta) \\ &\quad - q_1(0) - 2q_4(0) - q_5(\theta) \\ &= \frac{1}{4} \begin{pmatrix} -4\sin^2\theta & 0 & 0 & 0 \\ 0 & -5 - 2\cos\theta - 3\cos 2\theta + 2\cos 3\theta & 4\cos^2\frac{\theta}{2} & 2\sin\theta + 3\sin 2\theta - 2\sin 3\theta \\ 0 & 4\cos^2\frac{\theta}{2} & -2(3 + \cos 2\theta) & -2\sin\theta \\ 0 & 2\sin\theta + 3\sin 2\theta - 2\sin 3\theta & -2\sin\theta & 2\sin^2\frac{\theta}{2}\cos^2\frac{\theta}{2}(4\cos\theta - 3) \end{pmatrix}, \end{aligned} \quad (5)$$

again in the Bell basis and for quantum expressions similar to the ones enumerated in Eq. (1) [32].

In this basis, the operator $O_{33}(0, \theta, 2\theta, 0, \theta, 2\theta)$ splits into a direct sum of a one-dimensional part $-\sin^2\theta$ and a three-dimensional part o , respectively. Using the Cardano method (see Ref. [33]), one can solve the characteristic equation of the three-dimensional submatrix o

in the lower right corner of O_{33} ,

$$\lambda^3 + b(\theta)\lambda^2 + c(\theta)\lambda + d(\theta) = 0, \quad (6)$$

with the coefficients $b = -\text{Tr}o$, $c = 1/2(\text{Tr}^2o - \text{Tr}o^2)$, $d = -\text{det}o$. (For convenience we omit here the dependence on θ .) The (real) eigenvalues can then be written as [33]

$$\lambda_2 = -2\sqrt{|u|} \cos \frac{\xi}{3} - \frac{b}{3}, \quad (7)$$

$$\lambda_{3,4} = \sqrt{|u(x)|} \left[\cos \frac{\xi}{3} \pm \sin \frac{\xi}{3} \right] - \frac{b}{3},$$

with $u = 1/9(3c - b^2)$ and $\cos \xi = \frac{1}{54}(9bc - 2b^3 - 27d)/(u\sqrt{|u|})$. In Fig. 1, the eigenvalues $\lambda_2, \lambda_3, \lambda_4$, together with the eigenvalue $\lambda_1 = -\sin^2 \theta$ from the one-dimensional part of O_{33} , are plotted as functions of the parameter θ . The maximum violation of $1/4$ is obtained for $\theta = \pi/3$ with the associated eigenvector

$$|\Psi_{\max}\rangle = \frac{\sqrt{3}}{2} |\phi^-\rangle + \frac{1}{2} |\psi^+\rangle. \quad (8)$$

As indicated in Ref. [11], this scheme can be extended to m measurements on each particle by considering inequalities $I_{mm} \leq 0$ and corresponding operators O_{mm} of the form

$$I_{mm} = \sum_{j=1}^m \sum_{i=1}^{m-j+1} P(A_i B_j) - \sum_{i=1}^{m-1} P(A_{i+1} B_{m-i+1}) - \sum_{i=1}^m (m-i)P(B_i) - P(A_1) \leq 0, \quad (9)$$

where $P(A_i B_j)$ denotes the joint probability of obtaining the value one of the projection operators A_i and B_j operators on the left- and on the right-hand side, and $P(A_i), P(B_j)$ the marginal probabilities on one side, respectively. For a choice of measurement directions $\{0, \theta, 2\theta, \dots, m\theta\}$ on both sides, the maximizing eigenvalues are plotted in Fig. 2. The matrices belonging to the operators O_{mm} ($m \leq 6$) are of the same form as is O_{33} ; i.e., they split up into a direct sum of two matrices in the Bell basis; the maximal eigenvalues can therefore be calculated explicitly using Eqs. (6) and (7).

For experimental realizations of the O_{33} case and special parameter configurations, the *ansatz* of Cabello [22] and Bovino *et al.* [23] can be generalized to arbitrary local unitary transformations $U_{2 \times 2} \in \text{SU}(2) \otimes \text{SU}(2)$ ap-




FIG. 1. Eigenvalues of O_{33} in dependence of the relative angle θ .

plied to each one of the two particles in some Bell-basis state $|\varphi\rangle$ separately; i.e.,

$$U(\omega_1, \theta_1, \phi_1) \otimes U(\omega_2, \theta_2, \phi_2) |\varphi\rangle. \quad (10)$$

The single qubit operators are taken as $U(\omega, \theta, \phi) = e^{i\frac{\omega}{2}\vec{n}\cdot\vec{\sigma}} \in \text{SU}(2)$ with ω as the rotation angle about the axis $\vec{n} = (\sin\theta \cos\phi, \sin\theta \sin\phi, \cos\theta)^T$. For example, the use of the Bell state $|\psi^+\rangle$ and the successive application of the local unitary operation $U(\omega_1, \theta_1, \phi_1) \otimes U(\omega_2, \theta_2, \phi_2)$ with $\omega_1 = 2\pi/3, \theta_1 = \phi_1 = \pi/2$, and $\omega_2 = \theta_2 = \phi_2 = 0$ yields the maximally violating eigenvector $|\Psi_{\max}\rangle$ from Eq. (7) which is also maximally entangled.

For the general $m > 2$ case, however, it is not always possible to obtain all possible maximally violating states by starting from a Bell state: for general measurement angles, the experimental realization additionally requires a two-qubit transformation from $\text{SU}(4)/[\text{SU}(2) \otimes \text{SU}(2)]$, followed by a local unitary operation $U_{2 \times 2}$ in order to obtain all possible states [34]. As an example, consider the maximally violating but not maximally entangled state at $\theta = \pi/2$: $|\Psi_{\pi/2}\rangle \approx 0.86|\psi^+\rangle + 0.17|\psi^-\rangle + 0.47|\phi^-\rangle$ cannot be obtained from a Bell state, as entanglement is preserved under $\text{SU}(2) \otimes \text{SU}(2)$ operations.

Alternatively, multipoint interferometry [35–37] offers a direct proof-of-principle implementation: By choosing the appropriate transmission coefficients and phases in a generalized beam splitter setup, one can prepare any pure state from an input state $|11\rangle \equiv (0, 0, 0, 1)$ corresponding to a photon in a single input port. Take, for example, the maximal eigenstate of the O_{33} operator at $\theta = \pi/2$, $|\Psi_{\pi/2}\rangle \approx 0.86|\psi^+\rangle + 0.17|\psi^-\rangle + 0.47|\phi^-\rangle \equiv (0.34, 0.73, 0.49, -0.34)$. The appropriate transmission parameters can be calculated via the identification [35]

$$\begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}^T R(N)^{-1} = \begin{pmatrix} 0.34 \\ 0.73 \\ 0.49 \\ -0.34 \end{pmatrix}^T = \begin{pmatrix} e^{-i\phi_1} \cos \omega_1 \\ -e^{-i\phi_2} \cos \omega_2 \sin \omega_1 \\ e^{-i\phi_3} \cos \omega_3 \sin \omega_2 \sin \omega_1 \\ -\sin \omega_3 \sin \omega_2 \sin \omega_1 \end{pmatrix}^T \quad (11)$$

for $\omega_1 = 1.23, \omega_2 = 2.46, \omega_3 = 0.60$, and $\phi_1 = \phi_2 = \phi_3 = 0$, where $R(N)$ is a $\text{SU}(4)$ rotation serially composed by two-dimensional beam splitter matrices.

In summary, we have shown how to construct and experimentally test the exact quantum bounds of Bell-type inequalities by solving the eigenvalue problem of the associated self-adjoint transformation. Several problems remain open. Among them is the exact derivation of the quantum correlation hull [21,26], in particular, whether the quantum hull is obtainable by extending the classical Bell-type inequalities in the way as presented above; i.e.,




FIG. 2. Maximum violation of the operator O_{mm} for $m = 2, \dots, 6$ for a symmetric measurement setup; the longer dashes indicate larger m .

by substituting the quantum probabilities for the classical ones. This is by no means trivial, as the sections of the quantum hull need not necessarily be derivable by mere classical extensions. A second open question is related to the geometric structures arising from quantum expectation values. These need not necessarily be convex. Again, the question of direct extensibility remains open for the hull of quantum expectations from the classical ones.

This research has been supported by the Austrian Science Foundation (FWF), Project No. F1513. S.F. acknowledges helpful conversations with B. Hiesmayer and S. Scheel.

*Electronic address: sfilipp@ati.ac.at

†Electronic addresses: svozil@tuwien.ac.at
<http://tph.tuwien.ac.at/~svozil>

- [1] A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer Academic Publishers, Dordrecht, 1993).
- [2] B. S. Cirel'son, *Lett. Math. Phys.* **4**, 93 (1980).
- [3] B. S. Tsirelson, *Hadronic J. Suppl.* **8**, 329 (1993).
- [4] M. Froissart, *Nuovo Cimento Soc. Ital. Fis.* **64B**, 241 (1981).
- [5] I. Pitowsky, *J. Math. Phys. (N.Y.)* **27**, 1556 (1986).
- [6] I. Pitowsky, *Quantum Probability—Quantum Logic* (Springer, Berlin, 1989).
- [7] I. Pitowsky, in *Bell's Theorem, Quantum Theory and the Conceptions of the Universe*, edited by M. Kafatos (Kluwer, Dordrecht, 1989), pp. 37–49.
- [8] I. Pitowsky, *Math. Programming* **50**, 395 (1991).
- [9] I. Pitowsky, *Brit. J. Phil. Sci.* **45**, 95 (1994).

- [10] I. Pitowsky and K. Svozil, *Phys. Rev. A* **64**, 014102 (2001).
- [11] D. Collins and N. Gisin, [quant-ph/0306129](http://arxiv.org/abs/quant-ph/0306129).
- [12] C. Sliwa, *Phys. Lett. A* **317**, 165 (2003).
- [13] G. Boole, *An Investigation of the Laws of Thought* (Dover, New York, 1958).
- [14] G. Boole, *Philos. Trans. R. Soc. London* **152**, 225 (1862).
- [15] R. F. Werner and M. M. Wolf, *Phys. Rev. A* **64**, 032112 (2001).
- [16] G. Schachner [quant-ph/0312117](http://arxiv.org/abs/quant-ph/0312117).
- [17] B. S. Tsirel'son, *J. Sov. Math.* **36**, 557 (1987).
- [18] L. A. Khalfin and B. S. Tsirelson, *Found. Physics* **22**, 879 (1992).
- [19] A. Cabello, *Phys. Rev. Lett.* **88**, 060403 (2002).
- [20] A. Cabello, *Phys. Rev. A* **66**, 042114 (2002).
- [21] S. Filipp and K. Svozil, *Phys. Rev. A* **69**, 032101 (2004).
- [22] A. Cabello, *Phys. Rev. Lett.* **92**, 060403 (2004).
- [23] F. A. Bovino, G. Castagnoli, I. P. Degiovanni, and S. Castelletto, *Phys. Rev. Lett.* **92**, 060404 (2004).
- [24] P. R. Halmos, *Finite-Dimensional Vector Spaces* (Springer, New York, Heidelberg, Berlin, 1974).
- [25] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV: Analysis of Operators* (Academic Press, New York, 1978).
- [26] I. Pitowsky, in *Quantum Theory: Reconsideration of Foundations, Proceedings of the 2001 Växjö Conference* (World Scientific, Singapore, 2002), pp. 299–308.
- [27] Nondegenerate eigenstates are always representable by one-dimensional subspaces and thus are pure, the exception being the possibility of a mixing between degenerate eigenstates [28].
- [28] S. L. Braunstein, A. Mann, and M. Revzen, *Phys. Rev. Lett.* **68**, 3259 (1992).
- [29] Equivalent results hold for the Clauser-Horne-Shimony-Holt (CHSH) inequality [30,31].
- [30] D. N. Mermin, *Ann. N.Y. Acad. Sci.* **755**, 616 (1995).
- [31] J. L. Cereceda, *Found. Phys. Lett.* **14**, 401 (2001).
- [32] Note that q_1 refers to the state of the first particle, whereas q_4 and q_5 refer to the states of the second particle.
- [33] D. Cocolicchio and D. Viggiano, *J. Phys. A* **33**, 5669 (2000).
- [34] J. Zhang, J. Vala, S. Sastry, and K. B. Whaley, *Phys. Rev. A* **67**, 042313 (2003).
- [35] M. Reck, A. Zeilinger, H. J. Bernstein, and P. Bertani, *Phys. Rev. Lett.* **73**, 58 (1994).
- [36] M. Zukowski, A. Zeilinger, and M. A. Horne, *Phys. Rev. A* **55**, 2564 (1997).
- [37] K. Svozil, [quant-ph/0401113](http://arxiv.org/abs/quant-ph/0401113).