

A GLANCE AT SINGLET STATES AND FOUR-PARTITE CORRELATIONS

MARIA SCHIMPF — KARL SVOZIL

This paper is dedicated to Sylvia Pulmannová, in appreciation of her scientific accomplishments, and also for the enjoyment and privilege to be co-author in some of her research articles, as well as meeting her personally. One of the most remarkable impressions on me (K.S.) was her humbleness as a person and colleague, accompanied by her extreme professional skills in mathematics, logics and physics.

(Communicated by Anatolij Dvurečenskij)

ABSTRACT. Group theoretic methods to construct all N -particle singlet states by iterative recursion are presented. These techniques are applied to the quantum correlations of four spin one-half particles in their singlet states. Multipartite quantized systems can be partitioned, and their observables grouped and redefined into condensed correlations.

©2010
Mathematical Institute
Slovak Academy of Sciences

1. Introduction

Improved experimental particle production techniques and potential applications in quantum information theory have stimulated interest in multipartite singlet and other entangled states. In particular, singlet states are among the most useful states in quantum mechanics, as they appear form-invariant under spatial rotations. Hence, a physical property such as uniqueness [1] or equibalanced [2] which holds true in one frame or direction remains to be true in all other frames or directions obtained by spatial rotations.

Yet, the explicit structure of singlet states — although well understood in general terms in group theory — has up to now neither been enumerated nor investigated beyond a few instances for spin one-half and spin one particles. Recent theoretical and experimental studies in multi-particle production (e.g., [3])

2000 Mathematics Subject Classification: Primary 81P68.

Keywords: quantum information, singlet states, group theory, entanglement, quantum nonlocality.

suggest that a more systematic way to generate the complete set of arbitrary N -particle singlet states is desirable.

In the present study we first pursue an algorithmic generation strategy, and tabulate some of the first singlet states. The recursive method employed is based on triangle relations and Clebsch-Gordan coefficients (e.g., [4, Ch. 13, Sec. 27]). With this approach, a complete table of all angular momentum states can be enumerated. The singlet states are obtained *via* the various pathways towards the $j = m = 0$ states.

The procedure can best be illustrated by a triangular diagram, where the states in ascending order of angular momentum are drawn against the number of particles. In such a diagram, the “lowest” states correspond to singlet states.

Consider, for the sake of an explicit demonstration of this generation method, a two-dimensional diagram such as the ones depicted in Fig. 1 and in Fig. 2, which represents the “space” or “domain” of all multi-partite states. In such a diagram, the *number of particles* is represented by the abscissa (the x -coordinate) along the positive x -axis. The ordinate (the y -coordinate) of the state is equal the total angular momentum of the state. Note that a single point may represent many states; all corresponding to an equal number of particles, and all having the same total angular momentum. N -partite singlet states can be constructed by starting from the unique state of one particle, then proceeding *via* all “diagonal” and, whenever possible for integer spins, also “horizontal” pathways (e.g., the “horizontal” path in Fig. 2) consisting of single substeps adding one particle after the other — either diagonally from the lower left to the upper right “ \nearrow ”, or diagonally from the upper left to the lower right “ \searrow ”, or, if possible, also horizontally from left to right “ \rightarrow ” — towards the zero momentum state of N particles. Every diagonal or horizontal substep corresponds to the addition of a single particle. Below we shall explicitly construct singlet states composed from particles of spin one-half and spin one.

In the second part of this article, we present an explicit analysis of the singlet states of four spin one-half particles in terms of their probabilities and expectation functions for spin state measurements. We also investigate the possibility to group the outcomes of the four spin state measurements on each particle to obtain “condensed” observables. Likewise, we consider selection of one or two particles and the resulting correlations. One of our physical motivations for doing so was the question of how such “condensed” observables would perform with respect to violations of classical locality conditions.

2. General algorithm for obtaining singlet states

In what follows we present a method to construct all states for a given number of particles. They are the basis to construct non-trivial, e.g., non-“zigzag” singlet states, which are not just products of singlet states of a smaller number of particles. Although only the spin one-half and the spin one cases are explicitly discussed, the method applies to arbitrary spin.

2.1. Spin one-half

We start by considering the spin state of a single spin one-half particle. A second spin one-half particle is added by combining two angular momenta $\frac{1}{2}$ to all possible total angular momenta $l = 0$ or 1 . Next, a third particle is introduced by coupling a third angular momentum $\frac{1}{2}$ to all previously derived states. Following the triangle equation, the resulting j -values for each l are in the domain

$$|l - j_3| \leq j \leq l + j_3. \quad (1)$$

In order to obtain all N -particle singlet states, we successively produce all states (not only singlet states) of $N/2$ particles. Note that for $N/2 \leq h \leq N$ particles we only need angular momentum states within $0 \leq j \leq (N-h)/2$, because the construction method does not allow states with higher angular momentum to “bend diagonally backwards” and finally reach the angular momentum zero singlet state.

Angular momentum states will be written as $|h, j, m, i\rangle$, where h denotes the particle number, j the angular momentum, and m the magnetic quantum number. Note that there may exist many states with equal h , j and m . Thus i denotes the number of state in an enumeration of all h -partite states with identical angular momentum j and magnetic quantum number m . In the enumeration scheme chosen, we first take states generated from higher total angular momentum, followed by states with equal total angular momentum for spin one particles, and states with lower total angular momentum. For spin one-half particles, let us define a function $f(j+1, h)$ denoting the *total number* of states of h particles with total angular momentum $j/2$. This function is tabulated in Table 1 for the spin one-half particle case. The Clebsch-Gordan coefficient is denoted by $\langle j_1, j_2, m_1, m_2 | j, m \rangle$.

For spin one-half particles, an arbitrary state $|h, j, m, i\rangle$, $h > 1$, can be generated from the states with one particle less by *adding* a particle, thereby *increasing* or *decreasing* the total angular momentum of the previous state containing one particle less. Thus, we obtain two different pathways towards $|h, j, m, i\rangle$; one from the total angular momentum $j + \frac{1}{2}$, symbolized graphically by “ \searrow ”, and one from the total angular momentum $j - \frac{1}{2}$, symbolized graphically by “ \nearrow ”.

For the sake of demonstration of the method employed, we shall explicitly discuss one of the two cases, in which the *addition* of one particle $h - 1 \rightarrow h$ results in a *lowering* of the total angular momentum by $\frac{1}{2}$ through $j + \frac{1}{2} \rightarrow j$, thus representing the diagonal pathway “ \searrow ” from the “upper left” to the “lower right” in a diagram (nonuniquely) representing states as points with coordinates given by the number of particles and the total angular momentum, respectively (e.g., Fig. 1). The first contribution, associated with the magnetic quantum numbers $m - \frac{1}{2}$ and $+\frac{1}{2}$, can be constructed from the product state

$$|h - 1, j + \frac{1}{2}, m - \frac{1}{2}, i\rangle \otimes |1, \frac{1}{2}, \frac{1}{2}, 1\rangle \quad (2)$$

by multiplying it with the Clebsch-Gordan coefficient

$$\langle j + \frac{1}{2}, m - \frac{1}{2}, \frac{1}{2}, \frac{1}{2} | j, m \rangle. \quad (3)$$

Similarly, the second contribution to $|h, j, m, i\rangle$, associated with the magnetic quantum numbers $m + \frac{1}{2}$ and $-\frac{1}{2}$, can be constructed *via* the product state

$$|h - 1, j + \frac{1}{2}, m + \frac{1}{2}, i\rangle \otimes |1, \frac{1}{2}, -\frac{1}{2}, 1\rangle, \quad (4)$$

multiplied with the Clebsch-Gordan coefficient

$$\langle j + \frac{1}{2}, m + \frac{1}{2}, \frac{1}{2}, -\frac{1}{2} | j, m \rangle. \quad (5)$$

Adding the two results, we obtain the state $|h, j, m, i\rangle$; i.e.,

$$\begin{aligned} |h, j, m, i\rangle &= \langle j + \frac{1}{2}, m - \frac{1}{2}, \frac{1}{2}, \frac{1}{2} | j, m \rangle |h - 1, j + \frac{1}{2}, m - \frac{1}{2}, i\rangle \otimes |1, \frac{1}{2}, \frac{1}{2}, 1\rangle \\ &\quad + \langle j + \frac{1}{2}, m + \frac{1}{2}, \frac{1}{2}, -\frac{1}{2} | j, m \rangle |h - 1, j + \frac{1}{2}, m + \frac{1}{2}, i\rangle \otimes |1, \frac{1}{2}, -\frac{1}{2}, 1\rangle. \end{aligned} \quad (6)$$

We do this for $m = -j, \dots, j$ and for all states labeled by the state number $i = 1, 2, \dots, f((2j + 1) + 1, h - 1)$. Recall that $f(j + 1, h)$ denotes the total number of states of h particles with angular momentum $j/2$. It can be computed by counting the number of all states generated by all possible pathways in the construction method described above.

Similarly, if j is greater than zero, we obtain the state $|h, j, m, i\rangle$ from the diagonal pathway “ \nearrow ”, starting from the states $|h - 1, j - \frac{1}{2}, m - \frac{1}{2}, i\rangle$ and $|h - 1, j - \frac{1}{2}, m + \frac{1}{2}, i\rangle$ of $h - 1$ particles and total angular momentum $j - \frac{1}{2}$ by adding a single particle *via*

$$\langle j - \frac{1}{2}, m - \frac{1}{2}, \frac{1}{2}, \frac{1}{2} | j, m \rangle |h - 1, j - \frac{1}{2}, m - \frac{1}{2}, i\rangle \otimes |1, \frac{1}{2}, \frac{1}{2}, 1\rangle \quad (7)$$

and

$$\langle j - \frac{1}{2}, m + \frac{1}{2}, \frac{1}{2}, -\frac{1}{2} | j, m \rangle |h - 1, j - \frac{1}{2}, m + \frac{1}{2}, i\rangle \otimes |1, \frac{1}{2}, -\frac{1}{2}, 1\rangle. \quad (8)$$

This procedure is carried out for $m = -j, \dots, j$ and i satisfying

$$f((2j + 1) + 1, h - 1) + 1 \leq i \leq f((2j + 1) + 1, h - 1) + f((2j + 1) - 1, h - 1). \quad (9)$$

A concrete example is drawn in Fig. 1.

A GLANCE AT SINGLET STATES AND FOUR-PARTITE CORRELATIONS




FIGURE 1. Construction of both singlet states a) and b) of four spin one-half particles. Concentric circles indicate the target states. The second state is a “zigzag” state composed by the product of two two-partite singlet states

It contains the pathways leading to the construction of both singlet states of four spin one-half particles.

For spin one-half particles, the function $f(j + 1, h)$ denoting the *total number* of states of h particles with total angular momentum $j/2$ is tabulated in Table 1. The bottom line above the axis contains the number of different orthogonal singlet states.

The singlet states of up to six spin one-half particles are explicitly enumerated in Table 2.

2.2. Spin one

The construction of the singlet states of spin one particles follows similar rules as in the case of spin one-half particles. One example is the construction of the singlet state consisting of three spin one particles drawn in Fig. 2. Note that in this case, as for all particles of integer spin, there are *three* possible subpaths per addition of one particle; two diagonal “↗” and “↘” pathways as in the case for spin one-half particles, as well as one horizontal “→”.

Table 3 enumerates the numbers of states contributing to a calculation of singlet states up to 18 spin one particles. The bottom line above the axis shows the actual number of different orthogonal singlet states. The singlet states of up to four spin one (with one singlet state of 5) particles are explicitly enumerated in Table 4.

There always exist trivial “zigzag” singlet states which are the product of r two-particle singlet states stemming from the rising and lowering of consecutive

A GLANCE AT SINGLET STATES AND FOUR-PARTITE CORRELATIONS




FIGURE 2. Construction of the singlet state of three spin one particles. Note that for integer spin, there are three possible subpaths per addition of one particle; two diagonal “ \nearrow ” and “ \searrow ”, as well as one horizontal “ \rightarrow ”




FIGURE 3. Construction of the “zigzag” singlet state of N particles which effectively is a product state of $\frac{N}{2}$ spin l particle states

TABLE 2. First singlet states of N spin one-half particles

N	#	
2	1	$\frac{1}{\sqrt{2}}(+-\rangle - -+\rangle);$
4	1	$-\frac{1}{2\sqrt{3}}(-+-+\rangle + -++-\rangle + +- -+\rangle + +- +-\rangle) +$ $+\frac{1}{\sqrt{3}}(--++\rangle + ++--\rangle);$
4	2	$(-\frac{1}{\sqrt{2}} -+\rangle + \frac{1}{\sqrt{2}} +-\rangle)^2;$
6	1	$-\frac{1}{2} - - - + + +\rangle + -\frac{1}{6}(- + + - - +\rangle + - + + - + -\rangle +$ $+ - + + + - -\rangle + + - + - - +\rangle + + - + - + -\rangle +$ $+ + - + + - -\rangle + ++ - - - +\rangle + ++ - - + -\rangle +$ $+ ++ + - - -\rangle) + \frac{1}{6}(- - + - + +\rangle + - - + + - +\rangle +$ $+ - - + + + -\rangle + - + - - + +\rangle + - + - + - +\rangle +$ $+ - + - + + -\rangle + +- - - + +\rangle + +- - + - +\rangle +$ $+ +- - + + -\rangle) + \frac{1}{2} ++ + - - -\rangle;$
6	2	$-\frac{\sqrt{2}}{3} - - - + - +\rangle + -\frac{1}{3\sqrt{2}}(- + + + - -\rangle + +- + + - -\rangle +$ $+ ++ + - - -\rangle + ++ + - - +\rangle) + -\frac{1}{6\sqrt{2}}(- + - + - +\rangle +$ $+ - + - + + -\rangle + +- - + - +\rangle + +- - + + -\rangle) +$ $+\frac{1}{6\sqrt{2}}(- + + - - +\rangle + - + + - + -\rangle + +- + - - +\rangle +$ $+ +- + - + -\rangle) + \frac{1}{3\sqrt{2}}(- - + + - +\rangle + - - + + + -\rangle +$ $+ - + - - + +\rangle + +- - - + +\rangle) + \frac{\sqrt{2}}{3} ++ + - - -\rangle;$
6	3	$-\frac{1}{\sqrt{6}}(- + - - + +\rangle + - + + + - -\rangle) + -\frac{1}{2\sqrt{6}}(+ - - + - +\rangle +$ $+ + - - + + -\rangle + +- + - - +\rangle + +- + - + -\rangle) +$ $+\frac{1}{2\sqrt{6}}(- + - + - +\rangle + - + - + + -\rangle + - + + - - +\rangle +$ $+ - + + - + -\rangle) + \frac{1}{\sqrt{6}}(+ - - - + +\rangle + +- + + - -\rangle);$
6	4	$-\frac{1}{\sqrt{6}}(- - + + - +\rangle + ++ + - - +\rangle) + -\frac{1}{2\sqrt{6}}(- + - + + -\rangle +$ $+ - + + - + -\rangle + +- - + + -\rangle + +- + - + -\rangle) +$ $+\frac{1}{2\sqrt{6}}(- + - + - +\rangle + - + + - - +\rangle + +- - + - +\rangle +$ $+ +- + - - +\rangle) + \frac{1}{\sqrt{6}}(- - + + + -\rangle + ++ + - - +\rangle);$
6	5	$(-\frac{1}{\sqrt{2}} -+\rangle + \frac{1}{\sqrt{2}} +-\rangle)^3.$

3. Symmetries

In what follows we shall discuss the symmetry behavior of singlet states. In our approach the singlet states are orthogonal to each other. This can be

A GLANCE AT SINGLET STATES AND FOUR-PARTITE CORRELATIONS

TABLE 3. Enumeration of the total numbers of states contributing to a calculation of singlet states up to 18 spin one particles. The bottom line above the axis shows the actual number of different orthogonal singlet states

J	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	N		
0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
1	3	6	15	36	91	232	603	1585	4213	11298	30537	83097	227475	625992	1730787	4642	24882	102102	334425	1730787	
2	6	15	40	105	280	750	2025	5500	15026	41262	113841	315420	877320	227475	625992	1730787	4642	24882	102102	334425	1730787
3	10	29	84	238	672	1890	5313	14938	42042	118482	334425	877320	227475	625992	1730787	4642	24882	102102	334425	877320	1730787
4	15	49	154	468	1398	4125	12078	35178	102102	298500	830970	2274750	6259920	17307870	46420	248820	1021020	3344250	8773200	22747500	17307870
5	21	76	258	837	2640	8162	24882	75582	227475	707700	2274750	7077000	22747500	70770000	227475000	707700000	2274750000	7077000000	22747500000	70770000000	70770000000
6	28	111	405	1397	4642	15505	50520	163290	527280	1632900	5272800	16329000	52728000	163290000	527280000	1632900000	5272800000	16329000000	52728000000	163290000000	163290000000
7	36	155	605	24882	98019	381828	1488480	5818200	22747500	89019000	344425000	1330787000	5112980000	19633700000	75582000000	290102000000	1102102000000	4204200000000	15848200000000	59920000000000	227475000000000
8	45	2025	83097	334425	1330787	5112980	19633700	75582000	290102000	1102102000	4204200000	15848200000	59920000000	227475000000	890190000000	3444250000000	13307870000000	51129800000000	196337000000000	755820000000000	2901020000000000
9	55	2520	102102	4000000	15848200	62599200	248820000	980190000	3818280000	14884800000	58182000000	227475000000	890190000000	3444250000000	13307870000000	51129800000000	196337000000000	755820000000000	2901020000000000	11021020000000000	42042000000000000

TABLE 4. First singlet states of N spin one particles

N	#	
2	1	$\frac{1}{\sqrt{3}}(- 0,0\rangle + -1,1\rangle + 1,-1\rangle)$;
3	1	$-\frac{1}{\sqrt{6}}(-1,0,1\rangle + 0,1,-1\rangle + 1,-1,0\rangle)$ $+\frac{1}{\sqrt{6}}(-1,1,0\rangle + 0,-1,1\rangle + 1,0,-1\rangle)$;
4	1	$-\frac{1}{2\sqrt{5}}(-1,0,0,1\rangle + -1,0,1,0\rangle + 0,-1,0,1\rangle + 0,-1,1,0\rangle$ $+ 0,1,-1,0\rangle + 0,1,0,-1\rangle + 1,0,-1,0\rangle + 1,0,0,-1\rangle)$ $+\frac{1}{6\sqrt{5}}(-1,1,-1,1\rangle + -1,1,1,-1\rangle + 1,-1,-1,1\rangle$ $+ 1,-1,1,-1\rangle) + \frac{1}{3\sqrt{5}}(-1,1,0,0\rangle + 0,0,-1,1\rangle + 0,0,1,-1\rangle$ $+ 1,-1,0,0\rangle) + \frac{2}{3\sqrt{5}} 0,0,0,0\rangle + \frac{1}{\sqrt{5}}(-1,-1,1,1\rangle + 1,1,-1,-1\rangle)$;
4	2	$-\frac{1}{2\sqrt{3}}(-1,0,1,0\rangle + -1,1,-1,1\rangle + 0,-1,0,1\rangle + 0,1,0,-1\rangle$ $+ 1,-1,1,-1\rangle + 1,0,-1,0\rangle) + \frac{1}{2\sqrt{3}}(-1,0,0,1\rangle + -1,1,1,-1\rangle$ $+ 0,-1,1,0\rangle + 0,1,-1,0\rangle + 1,-1,-1,1\rangle + 1,0,0,-1\rangle)$;
4	3	$(\frac{1}{\sqrt{3}}(- 0,0\rangle + -1,1\rangle + 1,-1\rangle))^2$;
5	1	$-\sqrt{\frac{2}{15}} -1,-1,0,1,1\rangle + -\frac{1}{\sqrt{30}}(-1,0,1,0,0\rangle + 0,-1,1,0,0\rangle$ $+ 0,0,-1,0,1\rangle + 0,0,-1,1,0\rangle + 0,1,1,-1,-1\rangle$ $+ 1,0,1,-1,-1\rangle + 1,1,-1,-1,0\rangle + 1,1,-1,0,-1\rangle)$ $+ -\frac{1}{2\sqrt{30}}(-1,0,1,-1,1\rangle + -1,0,1,1,-1\rangle + -1,1,-1,0,1\rangle$ $+ -1,1,-1,1,0\rangle + 0,-1,1,-1,1\rangle + 0,-1,1,1,-1\rangle$ $+ 0,1,0,-1,0\rangle + 0,1,0,0,-1\rangle + 1,-1,-1,0,1\rangle$ $+ 1,-1,-1,1,0\rangle + 1,0,0,-1,0\rangle + 1,0,0,0,-1\rangle)$ $+\frac{1}{2\sqrt{30}}(-1,0,0,0,1\rangle + -1,0,0,1,0\rangle + -1,1,1,-1,0\rangle$ $+ -1,1,1,0,-1\rangle + 0,-1,0,0,1\rangle + 0,-1,0,1,0\rangle$ $+ 0,1,-1,-1,1\rangle + 0,1,-1,1,-1\rangle + 1,-1,1,-1,0\rangle$ $+ 1,-1,1,0,-1\rangle + 1,0,-1,-1,1\rangle + 1,0,-1,1,-1\rangle)$ $+\frac{1}{\sqrt{30}}(-1,-1,1,0,1\rangle + -1,-1,1,1,0\rangle + -1,0,-1,1,1\rangle$ $+ 0,-1,-1,1,1\rangle + 0,0,1,-1,0\rangle + 0,0,1,0,-1\rangle$ $+ 0,1,-1,0,0\rangle + 1,0,-1,0,0\rangle) + \sqrt{\frac{2}{15}} 1,1,0,-1,-1\rangle$;

demonstrated by considering the formula [5]

$$\begin{aligned}
 \langle (j'_1 j'_2) j m | (j_1 j_2) j m \rangle &= \sum_{m'_1+m'_2=m, m_1+m_2=m} \langle (j'_1 j'_2) j m | j'_1 m'_1 j'_2 m'_2 \rangle \times \\
 &\quad \times \langle j'_1 m'_1 j'_2 m'_2 | j_1 m_1 j_2 m_2 \rangle \langle j_1 m_1 j_2 m_2 | (j_1 j_2) j m \rangle \quad (10) \\
 &= \delta_{j_1 j'_1} \delta_{j_2 j'_2} \delta_{m_1 m'_1} \delta_{m_2 m'_2},
 \end{aligned}$$

where $|(j_1 j_2) j m\rangle$ stands for a state of total angular momentum j and magnetic quantum number m , composed of two parts having angular momentum j_1 and j_2 , respectively. States stemming from different j_1 values are orthogonal to each other. Hence, also the singlet states derived from them are orthogonal. By iteration it follows that even singlet states stemming from the same j_1 are orthogonal. The method allows us to construct the full basis for each singlet space which has the appropriate dimension.

3.1. Sign changes of magnetic quantum numbers

For the Clebsch-Gordan coefficients the following formula holds

$$\langle j_1, -m_1, j_2, -m_2 | j, -m \rangle = (-1)^{j_1+j_2-j} \langle j_1 m_1 j_2 m_2 | j m \rangle. \quad (11)$$

3.1.1. Spin one-half

In what follows, the symmetries of singlet spin one-half particle states are investigated. For a coupling j to $j + \frac{1}{2}$, the Clebsch-Gordan coefficients satisfy

$$\begin{aligned} \langle j, -m - \frac{1}{2}, \frac{1}{2}, \frac{1}{2} | j + \frac{1}{2}, -m \rangle &= (-1)^0 \langle j, m + \frac{1}{2}, \frac{1}{2}, -\frac{1}{2} | j + \frac{1}{2}, m \rangle \\ \langle j, m + \frac{1}{2}, \frac{1}{2}, -\frac{1}{2} | j + \frac{1}{2}, m \rangle &= (-1)^0 \langle j, -m - \frac{1}{2}, \frac{1}{2}, \frac{1}{2} | j + \frac{1}{2}, -m \rangle. \end{aligned}$$

If all the magnetic quantum numbers reverse their signs, the Clebsch-Gordan coefficients stay the same. Coupling $j + \frac{1}{2}$ to j results in

$$\begin{aligned} \langle j + \frac{1}{2}, m, \frac{1}{2}, \frac{1}{2} | j, m + \frac{1}{2} \rangle &= (-1)^1 \langle j + \frac{1}{2}, -m, \frac{1}{2}, -\frac{1}{2} | j, -m - \frac{1}{2} \rangle \\ \langle j + \frac{1}{2}, -m, \frac{1}{2}, -\frac{1}{2} | j, -m - \frac{1}{2} \rangle &= (-1)^1 \langle j + \frac{1}{2}, m, \frac{1}{2}, \frac{1}{2} | j, m + \frac{1}{2} \rangle. \end{aligned}$$

In this case, all the Clebsch-Gordan coefficients change their signs.

We conclude that the symmetry behavior remains the same if one passes from the angular momentum subspace $|N, J\rangle$ to the angular momentum subspace $|N + 1, J + \frac{1}{2}\rangle$. By passing from the subspace $|N, J\rangle$ to the subspace $|N + 1, J - \frac{1}{2}\rangle$ the symmetry behaviour changes from even to odd and from odd to even, respectively. A graphical representation of this property is depicted in Fig. 4. In particular, the singlet states where N is $k \cdot 2 \cdot 2$ (k is an integer) are even, and the ones where N is $2 \cdot (2k + 1)$ are odd.

3.1.2. Spin one

Let us now consider the $j = 1$ case first. For the coupling of j to $j + 1$, the symmetry described above implies

$$\begin{aligned} \langle j, -m - 1, 1, 1 | j + 1, -m \rangle &= (-1)^0 \langle j, m + 1, 1, -1 | j + 1, m \rangle, \\ \langle j, -m, 1, 0 | j + 1, -m \rangle &= (-1)^0 \langle j, m, 1, 0 | j + 1, m \rangle; \end{aligned} \quad (12)$$




FIGURE 4. Symmetry behavior of spin one-half particles. Even and odd subspaces are denoted by concentric and filled circles, respectively. The numbers denote the dimensions of the subspaces. The first number stands for the number of states $|h, j\rangle$, and the second number stands for the $2j + 1$ projections. Arrows represent the direction of the coupling

i.e., the Clebsch-Gordan coefficients are the same. For the coupling of j to j ,

$$\begin{aligned} \langle j, -m - 1, 1, 1 | j, -m \rangle &= (-1)^1 \langle j, m + 1, 1, -1 | j, m \rangle, \\ \langle j, -m, 1, 0 | j, -m \rangle &= (-1)^1 \langle j, m, 1, 0 | j, m \rangle; \end{aligned} \tag{13}$$

i.e., all Clebsch-Gordan coefficients change sign. Similarly for the coupling of $j + 1$ to j ,

$$\begin{aligned} \langle j + 1, m, 1, 1 | j, m + 1 \rangle &= (-1)^2 \langle j + 1, -m, 1, -1 | j, -m - 1 \rangle, \\ \langle j + 1, m, 1, 0 | j, m \rangle &= (-1)^2 \langle j + 1, -m, 1, 0 | j, -m \rangle; \end{aligned} \tag{14}$$

i.e., they all stay the same. Using these symmetries, we conclude that the symmetry behaviour remains the same if one passes from the angular momentum subspace $|N, j\rangle$ to the angular momentum subspace $|N + 1, j + 1\rangle$. The symmetry behaviour does not change for coupling $|N, j + 1\rangle$ to $|N + 1, j\rangle$. Coupling $|N, j\rangle$ to $|N + 1, j\rangle$ changes the symmetry behaviour from even to odd and from odd to even. The situation is depicted in Fig. 5. N -particle singlet states with N even are even, whereas N -particle singlet states with N odd are odd.

3.2. Symmetric group

Let us consider the permutations of the N magnetic quantum numbers in every product state of N particles. More explicitly, since every permutation




FIGURE 5. Symmetries of spin one particle states. Even subspaces are denoted by concentric circles, odd subspaces are denoted by filled circles. The numbers denote the dimensions of the subspaces. The first number stands for the number of states $|h, j\rangle$ and the second stands for the $2j + 1$ projections. Arrows represent the way of coupling

of N particles can be written as the product of $N - 1$ transpositions, we shall study the effects of $N - 1$ transpositions. We analyze $N - 1$ transpositions of the form $(j, j + 1)$, the transposition of j and $j + 1$ which generate the whole symmetric group, and in particular all the $N \cdot (N - 1)/2$ transpositions, since $(j, k + 1) = (k, k + 1)(j, k)(k, k + 1)$. Therefore, we consider the class $(2 \ 1^{N-2})$ of all two particle transpositions. Each irreducible representation can be labeled by an ordered partition of integers which corresponds to a specific Young diagram.

As stated in [4, App. D, Sec. 14], the space spanned by the vectors of total spins (SM) formed by N identical spins $\frac{1}{2}$ is associated with an irreducible representation of S_N , the representation whose Young diagram corresponds to the partition $[\frac{1}{2}N + S, \frac{1}{2}N - S]$ of the integer N . It is apparent that the Young diagrams for the irreducible components of the representation of S_N have at most two lines. For $N > 2$, any state contains at least two individual spins in the same state. Suppose the state contains the factor $u_+^{(i)}u_+^{(j)}$; i.e., $m_i, m_j = \frac{1}{2}$. Since $A = \frac{1}{2}(1 - (i, j))$ is the antisymmetrizer and $\frac{1}{2}(1 - (i, j))u_+^{(i)}u_+^{(j)} = 0$, it follows that $A|jm\rangle = 0$.

Using the theorem mentioned above, the Young diagrams of the irreducible spaces of the N -particle singlet states correspond to the partitions $[\frac{1}{2}N, \frac{1}{2}N]$. Hence the two-particle singlet state (sometimes referred to as the “Bell” state)

is an antisymmetric one-dimensional space. The four- and six-particle singlet spaces form a two- and a five-dimensional irreducible space whose Young diagrams are of the form $[2, 2]$ and $[3, 3]$. Using the formula for the dimension of an irreducible representation having the partition $[\lambda]$ (e.g., [6])

$$f^\lambda = n! \frac{\prod_{i < j \leq k} (\lambda_i - \lambda_j + j - i)}{\prod_{i=1}^k (\lambda_i + k - i)!}, \quad (15)$$

the dimension can be verified.

4. Four spin one-half particle correlations

Singlet states $|\Psi_{d,n,i}\rangle$ can be labeled by three numbers d , n and i , denoting the number d of outcomes associated with the dimension of Hilbert space per particle, the number n of participating particles, and the state count i in an enumeration of all possible singlet states of n particles of spin $j = (d - 1)/2$, respectively. To begin with the analysis of four-partite correlations, consider four spin one-half particles in one of the two singlet states enumerated in Table 2 and computed by following the “paths” indicated in Fig. 1; i.e.,

$$|\Psi_{2,4,1}\rangle = \frac{1}{\sqrt{3}} \left[|++--\rangle + |--++\rangle - \frac{1}{2}(|+-\rangle + |-+\rangle)(|+-\rangle + |-+\rangle) \right], \quad (16)$$

$$|\Psi_{2,4,2}\rangle = (|\Psi_{2,2,1}\rangle)^2 = \frac{1}{2}(|+-\rangle - |-+\rangle)(|+-\rangle - |-+\rangle), \quad (17)$$

where $|\Psi_{2,2,1}\rangle = \frac{1}{\sqrt{2}}(|+-\rangle - |-+\rangle)$ is the two particle singlet “Bell” state.

These pure states have an explicit vector space representation as orthogonal vectors. The two states corresponding to spin “up” and “down” correspond to $|+\rangle \equiv \hat{\mathbf{e}}_1 = (1, 0)$ and $|-\rangle \equiv \hat{\mathbf{e}}_2 = (0, 1)$. Product states can be represented by the tensor or Kronecker product, which, for two arbitrary vectors $\mathbf{a} = (a_1, a_2, \dots, a_n)$ and $\mathbf{b} = (b_1, b_2, \dots, b_m)$, can be represented by

$$\mathbf{a} \otimes \mathbf{b} = (a_1\mathbf{b}, a_2\mathbf{b}, \dots, a_n\mathbf{b}) = (a_1b_1, a_1b_2, \dots, a_nb_m). \quad (18)$$

Thus, by summing up all product states, the two singlet states have a vector representation as

$$\begin{aligned}\hat{\Psi}_{2,4,1} &= \frac{1}{\sqrt{3}} \left[\hat{\mathbf{e}}_1 \otimes \hat{\mathbf{e}}_1 \otimes \hat{\mathbf{e}}_2 \otimes \hat{\mathbf{e}}_2 + \hat{\mathbf{e}}_2 \otimes \hat{\mathbf{e}}_2 \otimes \hat{\mathbf{e}}_1 \otimes \hat{\mathbf{e}}_1 \right. \\ &\quad \left. - \frac{1}{\sqrt{2}} (\hat{\mathbf{e}}_1 \otimes \hat{\mathbf{e}}_2 + \hat{\mathbf{e}}_2 \otimes \hat{\mathbf{e}}_1) \otimes \frac{1}{\sqrt{2}} (\hat{\mathbf{e}}_1 \otimes \hat{\mathbf{e}}_2 + \hat{\mathbf{e}}_2 \otimes \hat{\mathbf{e}}_1) \right] \\ &= \left(0, 0, 0, \frac{1}{\sqrt{3}}, 0, -\frac{1}{2\sqrt{3}}, -\frac{1}{2\sqrt{3}}, 0, 0, -\frac{1}{2\sqrt{3}}, -\frac{1}{2\sqrt{3}}, 0, \frac{1}{\sqrt{3}}, 0, 0, 0 \right). \quad (19)\end{aligned}$$

$$\begin{aligned}\hat{\Psi}_{2,4,2} &= \frac{1}{\sqrt{2}} (\hat{\mathbf{e}}_1 \otimes \hat{\mathbf{e}}_2 - \hat{\mathbf{e}}_2 \otimes \hat{\mathbf{e}}_1) \otimes \frac{1}{\sqrt{2}} (\hat{\mathbf{e}}_1 \otimes \hat{\mathbf{e}}_2 - \hat{\mathbf{e}}_2 \otimes \hat{\mathbf{e}}_1) \\ &= \left(0, 0, 0, 0, 0, \frac{1}{2}, -\frac{1}{2}, 0, 0, -\frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0 \right). \quad (20)\end{aligned}$$

Their density operators ρ_i , $i = 1, 2$, are just the projectors corresponding to the one-dimensional linear subspaces spanned by the vectors representing $\hat{\Psi}_{2,4,2}$ and $\hat{\Psi}_{2,4,1}$ in Eqs. (20, 19); i. e. they are the dyadic product

$$\rho_i = \left[\hat{\Psi}_{2,4,i}^T \hat{\Psi}_{2,4,i} \right]. \quad (21)$$

As has been pointed out above, and as $\hat{\Psi}_{2,4,2} \cdot \hat{\Psi}_{2,4,1} = 0$ or equivalently $\rho_{\Psi_{2,4,1}} \cdot \rho_{\Psi_{2,4,2}} = 0$, the singlet states are orthogonal. The most general form of a four spin one-half particle singlet state is thus given by

$$|\Psi_{2,4,s}\rangle = \lambda_1 |\Psi_{2,4,1}\rangle + \lambda_2 |\Psi_{2,4,2}\rangle \quad (22)$$

with $|\lambda_1|^2 + |\lambda_2|^2 = 1$, which can be parameterized by $\lambda_1 = \sin \tau$, $\lambda_2 = \cos \tau$, such that for $\tau = 0$, $|\Psi_{2,4,s}\rangle = |\Psi_{2,4,2}\rangle$, and for $\tau = \pi/2$, $|\Psi_{2,4,s}\rangle = |\Psi_{2,4,1}\rangle$.

Singlet states are form invariant with respect to arbitrary unitary transformations in the single-particle Hilbert spaces and thus also rotationally invariant in configuration space, in particular under the rotations $|+\rangle = e^{i\frac{\varphi}{2}} \left(\cos \frac{\theta}{2} |+\prime\rangle - \sin \frac{\theta}{2} |-\prime\rangle \right)$ and $|-\rangle = e^{-i\frac{\varphi}{2}} \left(\sin \frac{\theta}{2} |+\prime\rangle + \cos \frac{\theta}{2} |-\prime\rangle \right)$ in the spherical coordinates θ, φ defined below (e.g., [7, Eq. (2)], or [8, Eq. (7-49)]). However, despite this form invariance under rotations, the states are non-unique in the sense that knowledge of a spin state observable for one particle is not sufficient for the simultaneous (counterfactual) determination of spin state properties for all other three particles [9, 10].

4.1. Operators

In what follows, the operators corresponding to the spin state observables will be enumerated. Thereby, spherical coordinates represent angles of spin state measurements. Suppose that i denotes the i th particle with $1 \leq i \leq 4$. Let θ_i be the polar angle in the x - z -plane from the z -axis with $0 \leq \theta_i \leq \pi$, and φ_i the azimuthal angle in the x - y -plane from the x -axis with $0 \leq \varphi_i < 2\pi$. For the

sake of simplicity, we shall sometimes consider measurements in the x - z -plane, for which $\varphi_1 = \varphi_2 = \varphi_3 = \varphi_4 = 0$. Because of the spherical symmetry of the singlet state, this is in every aspect equivalent to a measurement along angles lying in an arbitrary plane. In such cases the expectation values (the raw, or uncentered, product moments [11]) are merely functions of the polar angles $\theta_1, \theta_2, \theta_3$ and θ_4 , so the azimuthal angles will be omitted. For compact notation, $\hat{\theta}$ and $\hat{\varphi}$ will be used to denote the coordinates $\theta_1, \theta_2, \theta_3, \theta_4$ and $\varphi_1, \varphi_2, \varphi_3, \varphi_4$, respectively.

The projection operators F corresponding to a four spin one-half particle joint measurement aligned (“+”) or antialigned (“-”) along those angles are

$$F_{\pm\pm\pm\pm}(\hat{\theta}, \hat{\varphi}) = \frac{1}{2} [\mathbb{I}_2 \pm \sigma(\theta_1, \varphi_1)] \otimes \frac{1}{2} [\mathbb{I}_2 \pm \sigma(\theta_2, \varphi_2)] \otimes \frac{1}{2} [\mathbb{I}_2 \pm \sigma(\theta_3, \varphi_3)] \otimes \frac{1}{2} [\mathbb{I}_2 \pm \sigma(\theta_4, \varphi_4)],$$

with $\sigma(\theta, \varphi) = \begin{pmatrix} \cos \theta & e^{-i\varphi} \sin \theta \\ e^{i\varphi} \sin \theta & -\cos \theta \end{pmatrix}$. For example, $F_{-+-+}(\hat{\theta}, \hat{\varphi})$ stands for the proposition

‘The spin state of the first particle measured along θ_1, φ_1 is “-”, the spin state of the second particle measured along θ_2, φ_2 is “+”, the spin state of the third particle measured along θ_3, φ_3 is “-”, and the spin state of the fourth particle measured along θ_4, φ_4 is “+”.’

Fig. 6 depicts a measurement configuration for a simultaneous measurement of spins along $\theta_1, \varphi_1, \theta_2, \varphi_2, \theta_3, \varphi_3$ and θ_4, φ_4 of the state $\Psi_{2,4,2}$.

4.2. Probabilities and expectations

We now turn to the calculation of quantum predictions. The joint probability to register the spins of the four particles in state $\rho_{\Psi_{2,4,s}}$ aligned or antialigned along the directions defined by $(\theta_1, \varphi_1), (\theta_2, \varphi_2), (\theta_3, \varphi_3)$, and (θ_4, φ_4) can be evaluated by a straightforward calculation of

$$P_{\rho_{\Psi_{2,4,s}}} \pm\pm\pm\pm(\hat{\theta}, \hat{\varphi}) = \text{Tr}[\rho_{\Psi_{2,4,s}} \cdot F_{\pm\pm\pm\pm}(\hat{\theta}, \hat{\varphi})]. \tag{23}$$

The expectation functions and joint probabilities to find the four particles in an even or in an odd number of spin “-”-states when measured along $(\theta_1, \varphi_1), (\theta_2, \varphi_2), (\theta_3, \varphi_3)$, and (θ_4, φ_4) are enumerated in Table 5. In the following, omitted arguments are zero. For example, the expectation function of the general singlet state in Eq. (22) restricted to $\varphi_1 = \varphi_2 = \varphi_3 = \varphi_4 = 0$ is

$$E(\tau; \hat{\theta}) = \frac{1}{3} \left([2 + \cos(2\tau)] \cos(\theta_1 - \theta_2) \cos(\theta_3 - \theta_4) + 2 \sin \tau [\sin \tau \cos(\theta_1 + \theta_2 - \theta_3 - \theta_4) + \sqrt{3} \cos \tau \sin(\theta_1 - \theta_2) \sin(\theta_3 - \theta_4)] \right) \tag{24}$$

A GLANCE AT SINGLET STATES AND FOUR-PARTITE CORRELATIONS




FIGURE 6. Simultaneous spin measurement of the four-partite singlet state represented in Eq. (17). Boxes indicate spin state analyzers such as Stern-Gerlach apparatus oriented along the directions θ_1, φ_1 , θ_2, φ_2 , θ_3, φ_3 and θ_4, φ_4 ; their two output ports are occupied with detectors associated with the outcomes “+” and “-”, respectively

For $\tau = 0$ and $\tau = \frac{\pi}{2}$, Eq. (24) reduces to $E_{\rho_{\Psi_{2,4,2}}}$ and $E_{\rho_{\Psi_{2,4,1}}}$ in Table 5, respectively.

We concentrate on the algebraic evaluation of $E_{\rho_{\Psi_{2,4,1}}}$, as this expectation function is from a nontrivial non-zigzag singlet state and thus can be expected to reveal additional structure not inherited from the two-partite correlations also enumerated in Table 5. If all the polar angles $\hat{\theta}$ are all set to $\pi/2$, then this correlation function yields

$$E_{\rho_{\Psi_{2,4,1}}}\left(\frac{\pi}{2}, \frac{\pi}{2}, \frac{\pi}{2}, \frac{\pi}{2}, \hat{\varphi}\right) = \frac{1}{3} [2 \cos(\varphi_1 + \varphi_2 - \varphi_3 - \varphi_4) + \cos(\varphi_1 - \varphi_2) \cos(\varphi_3 - \varphi_4)]. \quad (25)$$

Likewise, if all the azimuthal angles $\hat{\varphi}$ are all set to zero, one obtains

$$E_{\rho_{\Psi_{2,4,1}}}(\hat{\theta}) = \frac{1}{3} [2 \cos(\theta_1 + \theta_2 - \theta_3 - \theta_4) + \cos(\theta_1 - \theta_2) \cos(\theta_3 - \theta_4)]. \quad (26)$$

4.3. Plasticity of expectation function

The plasticity of the expectation function $E(\tau; \hat{\theta})$ is comparable to the two-particle expectation function $E(\theta) = -\cos\theta$ for measurements in one plane can be demonstrated by plotting the probabilities and expectation values for selectively chosen parameters, as depicted in Fig. 7.

As there are four particles involved, the outcomes of one or two particles can be utilized to select the events of the other particles. Let “ \pm_i ” stand for the

TABLE 5. Probabilities and expectation functions for finding an odd or even number of spin “-”-states. Omitted arguments are zero

Two-partite singlet state	
	$P_{=} = \frac{1}{2}(1 + E), \quad P_{\neq} = \frac{1}{2}(1 - E)$
	$E(\theta_1, \theta_2, \varphi_1, \varphi_2) = P_{=} - P_{\neq} = -[\cos \theta_1 \cos \theta_2 + \cos(\varphi_1 - \varphi_2) \sin \theta_1 \sin \theta_2]$
	$E(\theta_1, \theta_2) = -\cos(\theta_1 - \theta_2)$
	$E(\frac{\pi}{2}, \frac{\pi}{2}, \varphi_1, \varphi_2) = -\cos(\varphi_1 - \varphi_2)$
Four-partite singlet states	
	$P_{\text{even}} = \frac{1}{2}[1 + E], \quad P_{\text{odd}} = \frac{1}{2}[1 - E], \quad E = P_{\text{even}} - P_{\text{odd}}$
	$E_{\rho_{\Psi_{2,4,1}}}(\hat{\theta}, \hat{\varphi}) = \frac{1}{3} \{ \cos \theta_3 \sin \theta_1 [-\cos \theta_4 \cos(\varphi_1 - \varphi_2) \sin \theta_2$
	$+ 2 \cos \theta_2 \cos(\varphi_1 - \varphi_4) \sin \theta_4]$
	$+ \sin \theta_1 \sin \theta_3 [2 \cos \theta_2 \cos \theta_4 \cos(\varphi_1 - \varphi_3)$
	$+ (2 \cos(\varphi_1 + \varphi_2 - \varphi_3 - \varphi_4)$
	$+ \cos(\varphi_1 - \varphi_2) \cos(\varphi_3 - \varphi_4)] \sin \theta_2 \sin \theta_4]$
	$+ \cos \theta_1 [2 \sin \theta_2 (\cos \theta_4 \cos(\varphi_2 - \varphi_3) \sin \theta_3$
	$+ \cos \theta_3 \cos(\varphi_2 - \varphi_4) \sin \theta_4]$
	$+ \cos \theta_2 (3 \cos \theta_3 \cos \theta_4 - \cos(\varphi_3 - \varphi_4) \sin \theta_3 \sin \theta_4)] \}$
	$E_{\rho_{\Psi_{2,4,1}}}(\frac{\pi}{2}, \frac{\pi}{2}, \frac{\pi}{2}, \frac{\pi}{2}, \hat{\varphi}) = \frac{1}{3} [2 \cos(\varphi_1 + \varphi_2 - \varphi_3 - \varphi_4) + \cos(\varphi_1 - \varphi_2) \cos(\varphi_3 - \varphi_4)]$
	$E_{\rho_{\Psi_{2,4,1}}}(\hat{\theta}) = \frac{1}{3} [2 \cos(\theta_1 + \theta_2 - \theta_3 - \theta_4) + \cos(\theta_1 - \theta_2) \cos(\theta_3 - \theta_4)]$
	$E_{\rho_{\Psi_{2,4,2}}}(\hat{\theta}) = \cos(\theta_1 - \theta_2) \cos(\theta_3 - \theta_4)$
	$E_{\rho_{\Psi_{2,4,2}}}(\hat{\theta}, \hat{\varphi}) = [\cos \theta_1 \cos \theta_2 + \cos(\varphi_1 - \varphi_2) \sin \theta_1 \sin \theta_2] \cdot$
	$\cdot [\cos \theta_3 \cos \theta_4 + \cos(\varphi_3 - \varphi_4) \sin \theta_3 \sin \theta_4]$
	$E(\tau; \hat{\theta}) = \frac{1}{3} \{ [2 + \cos(2\tau)] \cos(\theta_1 - \theta_2) \cos(\theta_3 - \theta_4)$
	$+ 2 \sin \tau [\sin \tau \cos(\theta_1 + \theta_2 - \theta_3 - \theta_4)$
	$+ \sqrt{3} \cos \tau \sin(\theta_1 - \theta_2) \sin(\theta_3 - \theta_4)] \}$
	$E(\tau; \hat{\theta}, \hat{\varphi}) = \frac{1}{3} \{ \cos \theta_1 (\cos \theta_2 \{ 3 \cos \theta_3 \cos \theta_4$
	$+ [2 \cos(2\tau) + 1] \cos(\varphi_3 - \varphi_4) \sin \theta_3 \sin \theta_4 \}$
	$+ 2 \sin \theta_2 \sin \tau [\cos \theta_3 \cos(\varphi_2 - \varphi_4) \sin \theta_4 (\sqrt{3} \cos \tau + \sin \tau)$
	$- \cos \theta_4 \cos(\varphi_2 - \varphi_3) \sin \theta_3 (\sqrt{3} \cos \tau - \sin \tau)] \}$
	$+ \sin \theta_1 (\cos \theta_3 \{ \cos \theta_4 [2 \cos(2\tau) + 1] \cos(\varphi_1 - \varphi_2) \sin \theta_2$
	$+ 2 \cos \theta_2 \cos(\varphi_1 - \varphi_4) \sin \theta_4 \sin \tau (\sin \tau - \sqrt{3} \cos \tau) \}$
	$+ \sin \theta_3 [2 \cos \theta_2 \cos \theta_4 \cos(\varphi_1 - \varphi_3) \sin \tau (\sqrt{3} \cos \tau + \sin \tau)$
	$+ \sin \theta_2 \sin \theta_4 \{ 2 \cos(\varphi_1 + \varphi_2 - \varphi_3 - \varphi_4) \sin^2 \tau$
	$+ [\cos(2\tau) + 2] \cos(\varphi_1 - \varphi_2) \cos(\varphi_3 - \varphi_4)$
	$+ \sqrt{3} \sin(2\tau) \sin(\varphi_1 - \varphi_2) \sin(\varphi_3 - \varphi_4)] \}$

A GLANCE AT SINGLET STATES AND FOUR-PARTITE CORRELATIONS




FIGURE 7. Probabilities and expectation values for (a) $\tau = 0$, $\theta_1 = \theta$, $\theta_2 = \theta_3 = \theta_4 = 0$, (b) $\tau = 0$, $\theta_1 = \theta$, $\theta_2 = \theta_3 = 0$, $\theta_4 = \pi$, (c) $\tau = \frac{\pi}{2}$, $\theta_1 = \theta_2 = -\theta_3 = \theta_4 = \theta$, (d) $\tau = \frac{\pi}{2}$, $\theta_1 = -\theta_3 = \theta_4 = \theta$, $\theta_2 = \frac{\pi}{4}$, (e) $\tau = \frac{\pi}{4}$, $\theta_1 = -\theta_3 = \theta_4 = \theta$, $\theta_2 = \frac{\pi}{4}$, (f) $\tau = \frac{\pi}{4}$, $\theta_1 = -\theta_3 = \theta_4 = \theta$, $\theta_2 = 0$. Dashed (dash dotted) lines indicate probabilities to find an even (odd) number of “-” outcomes, solid lines depict expectation functions

TABLE 6. Probabilities and expectation functions for finding an odd or even number of spin “-”-states with selection. “ \pm_i ” stands for the observation of spin state plus or minus on the i th particle

Three-partite GHZM state ([7])	
$P_{\pm} = \frac{1}{4} [1 + 2E], \quad P_{\pm} = \frac{1}{4} [1 - 2E], \quad E_{\pm} = P_{\pm} - P_{\pm}$	
$E_{\pm}(\theta_1, \theta_2, \theta_3, \varphi_1, \varphi_2, \varphi_3 \pm_3) = \frac{1}{2} [\cos \theta_1 \cos \theta_2 \pm_3 \cos(\varphi_1 + \varphi_2 + \varphi_3) \sin \theta_1 \sin \theta_2 \sin \theta_3]$	
Four-partite singlet states	
$E_{\rho_{\Psi_{2,4,1}}}(\hat{\theta}, \hat{\varphi} \pm_4) = \frac{1}{12} \pm \frac{1}{2} E_{\rho_{\Psi_{2,4,1}}}(\hat{\theta}, \hat{\varphi})$	
$E_{\rho_{\Psi_{2,4,1}}}(\hat{\theta} \pm_3 \pm_4) = \frac{1}{12} \{ 2(\pm_3 1)(\pm_4 1) \cos(\theta_1 + \theta_2 - \theta_3 - \theta_4) + \cos(\theta_1 - \theta_2) [1 + (\pm_3 1)(\pm_4 1) \cos(\theta_3 - \theta_4)] \}$	
$E_{\rho_{\Psi_{2,4,1}}}(\hat{\theta}, \hat{\varphi} \pm_3 \pm_4) = \frac{1}{12} \{ \cos \theta_1 (2(\pm_3 1)(\pm_4 1) \sin \theta_2 [\cos \theta_4 \cos(\varphi_2 - \varphi_3) \sin \theta_3 + \cos \theta_3 \cos(\varphi_2 - \varphi_4) \sin \theta_4] + \cos \theta_2 [1 + 3(\pm_3 1)(\pm_4 1) \cos \theta_3 \cos \theta_4 - (\pm_3 1)(\pm_4 1) \cos(\varphi_3 - \varphi_4) \sin \theta_3 \sin \theta_4] + \sin \theta_1 (\cos(\varphi_1 - \varphi_2) \sin \theta_2 [1 - (\pm_3 1)(\pm_4 1) \cos \theta_3 \cos \theta_4 + (\pm_3 1)(\pm_4 1) \cos(\varphi_3 - \varphi_4) \sin \theta_3 \sin \theta_4] + 2(\pm_3 1)(\pm_4 1) [\cos \theta_2 \cos \theta_4 \cos(\varphi_1 - \varphi_3) \sin \theta_3 + \cos \theta_2 \cos \theta_3 \cos(\varphi_1 - \varphi_4) \sin \theta_4 + \cos(\varphi_1 + \varphi_2 - \varphi_3 - \varphi_4) \sin \theta_2 \sin \theta_3 \sin \theta_4]) \}$	
$E_{\rho_{\Psi_{2,4,1}}}(\hat{\theta} \pm_2 \pm_4) = \frac{1}{12} \{ (\pm_2 1)(\pm_4 1) [2 \cos(\theta_1 + \theta_2 - \theta_3 - \theta_4) + \cos(\theta_1 - \theta_2) \cos(\theta_3 - \theta_4)] - 2 \cos(\theta_1 - \theta_3) \}$	
$E_{\rho_{\Psi_{2,4,1}}}(\hat{\theta}, \hat{\varphi} \pm_2 \pm_4) = \frac{1}{12} \{ \cos \theta_1 ((\pm_2 1)(\pm_4 1) \sin \theta_3 [2 \cos \theta_4 \cos(\varphi_2 - \varphi_3) \sin \theta_2 - \cos \theta_2 \cos(\varphi_3 - \varphi_4) \sin \theta_4] + \cos \theta_3 [-2 + 3(\pm_2 1)(\pm_4 1) \cos \theta_2 \cos \theta_4 + 2(\pm_2 1)(\pm_4 1) \cos(\varphi_2 - \varphi_4) \sin \theta_2 \sin \theta_4]) + \sin \theta_1 ((\pm_2 1)(\pm_4 1) \cos \theta_3 [-\cos \theta_4 \cos(\varphi_1 - \varphi_2) \sin \theta_2 + 2 \cos \theta_2 \cos(\varphi_1 - \varphi_4) \sin \theta_4] + \sin \theta_3 [2 [-1 + (\pm_2 1)(\pm_4 1) \cos \theta_2 \cos \theta_4] \cos(\varphi_1 - \varphi_3) + (\pm_2 1)(\pm_4 1) (2 \cos(\varphi_1 + \varphi_2 - \varphi_3 - \varphi_4) + \cos(\varphi_1 - \varphi_2) \cos(\varphi_3 - \varphi_4)) \sin \theta_2 \sin \theta_4]) \}$	

observation of spin state plus or minus on the i th particle. Table 6 contains the results of the associated expectation values and joint probabilities for finding an odd or even number of spin “-”-states.

Two or three observables could also be grouped together to form a “condensed” observable. For instance, for each individual quadruple of outcomes

$\{o_1, o_2, o_3, o_4\}$ the values of the first and the second, as well as of the third and the fourth particle could be multiplied to obtain two other, dichotomic observables $o_1 o_2$ and $o_3 o_4$, respectively. More generally, one could take all partitions of 4, such that the outcomes of all particles within an element of the partition are multiplied. As the single outcomes occur at random, their resulting products and thus the new condensed observable would also represent random variables. Since the multiplication is associative, the resulting condensed correlations are just the four-partite correlations discussed so far.

5. Summary

In summary, we have discussed an algorithmic procedure to enumerate all singlet states of N particles of arbitrary spin. We have then explicitly enumerated the first cases for spin one-half and spin one and discussed their symmetries. These results have then be applied for a calculation of the quantum probabilities and expectation functions of four spin one-half particles in four arbitrary directions. We conclude by pointing out that all discussed configurations could, as a proof of principle, be locally realized by generalized beam splitters [12, 13, 14].

Acknowledgement. The authors thank Rainer Dirl and Peter Kasperkovitz for their advise in group theoretical questions; K.S. gratefully acknowledges discussions with Boris Kamenik, Günther Krenn and Johann Summhammer in Viennese coffee houses and elsewhere.

REFERENCES

- [1] SVOZIL, K.: *Are simultaneous Bell measurements possible?*, New J. Phys. **8** (2006), 39.
<http://dx.doi.org/10.1088/1367-2630/8/3/039>
- [2] ZEILINGER, A.: *A foundational principle for quantum mechanics*, Found. Phys. **29** (1999), 631–643.
<http://dx.doi.org/10.1023/A:1018820410908>
- [3] EIBL, M. — GAERTNER, S. — BOURENNANE, M. — KURTSIEFER, C. — ZUKOWSKI, M. — WEINFURTER, H.: *Experimental observation of four-photon entanglement from parametric down-conversion*, Phys. Rev. Lett. **90** (2003), 200403-1–200403-4.
<http://dx.doi.org/10.1103/PhysRevLett.90.200403>
- [4] MESSIAH, A.: *Quantum Mechanics. Vol. II*, North-Holland, Amsterdam, 1962.
- [5] CHAICHIAN, M.—HAGEDORN, R.: *Symmetries in Quantum Mechanics. From Angular Momentum to Supersymmetry*, Grand. Student Ser. Phys., Institute of Physics Publ., Bristol-Philadelphia, 1998.
- [6] WYBOURNE, B. G.: *Symmetry Principles and Atomic Spectroscopy*. Wiley Interscience, New York, 1970.

- [7] KRENN, G.—ZEILINGER, A.: *Entangled entanglement*, Phys. Rev. A (3) **54** (1996), 1793–1797.
<http://dx.doi.org/10.1103/PhysRevA.54.1793>
- [8] BALLENTINE, L. E.: *Quantum Mechanics*. Prentice Hall, Englewood Cliffs, NY, 1989.
- [9] EINSTEIN, A.—PODOLSKY, B.—ROSEN, N.: *Can quantum-mechanical description of physical reality be considered complete?*, Phys. Rev. **47** (1935), 777–780.
<http://dx.doi.org/10.1103/PhysRev.47.777>
- [10] SVOZIL, K.: *On counterfactuals and contextuality*. In: AIP Conference Proceedings 750 (Sweden, 2004). Foundations of Probability and Physics 3 (A. Khrennikov, ed.), American Institute of Physics, Melville, NY, 2005, pp. 351–360.
<http://dx.doi.org/10.1063/1.1874586>
- [11] GILL, R. D.: *Time, finite statistics, and Bell's fifth position*. In: Proceedings of Foundations of Probability and Physics 2 (Växjö, 2002) (A. Khrennikov, ed.), Växjö Univ. Press, Växjö, 2003, pp. 179–206.
- [12] RECK, M.—ZEILINGER, A.—BERNSTEIN, H. J.—BERTANI, P.: *Experimental realization of any discrete unitary operator*, Phys. Rev. Lett. **73** (1994), 58–61.
<http://dx.doi.org/10.1103/PhysRevLett.73.58>
- [13] ZUKOWSKI, M.—ZEILINGER, A.—HORNE, M. A.: *Realizable higher-dimensional two-particle entanglements via multipoint beam splitters*, Phys. Rev. A (3) **55** (1997), 2564–2579.
<http://dx.doi.org/10.1103/PhysRevA.55.2564>
- [14] SVOZIL, K.: *Noncontextuality in multipartite entanglement*, Amer. J. Phys. **38** (2005), 5781–5798.
<http://dx.doi.org/10.1088/0305-4470/38/25/013>

Received 12. 1. 2009
 Accepted 29. 1. 2009

*Institute for Theoretical Physics
 Vienna University of Technology
 Wiedner Hauptstraße 8-10/136
 A-1040 Vienna
 AUSTRIA
 E-mail: svozil@tuwien.ac.at
 URL: <http://tph.tuwien.ac.at/svozil>*